

Multigraded Castelnuovo-Mumford regularity

By *Diane Maclagan* at Stanford and *Gregory G. Smith* at New York

Abstract. We develop a multigraded variant of Castelnuovo-Mumford regularity. Motivated by toric geometry, we work with modules over a polynomial ring graded by a finitely generated abelian group. As in the standard graded case, our definition of multigraded regularity involves the vanishing of graded components of local cohomology. We establish the key properties of regularity: its connection with the minimal generators of a module and its behavior in exact sequences. For an ideal sheaf on a simplicial toric variety X , we prove that its multigraded regularity bounds the equations that cut out the associated subvariety. We also provide a criterion for testing if an ample line bundle on X gives a projectively normal embedding.

1. Introduction

Castelnuovo-Mumford regularity is a fundamental invariant in commutative algebra and algebraic geometry. Intuitively, it measures the complexity of a module or sheaf. The regularity of a module approximates the largest degree of the minimal generators and the regularity of a sheaf estimates the smallest twist for which the sheaf is generated by its global sections. Although the precise definition may seem rather technical, its value in bounding the degree of syzygies [20], [13] and constructing Hilbert schemes [18], [26] has established that regularity is an indispensable tool in both fields.

The goal of this paper is to develop a multigraded variant of Castelnuovo-Mumford regularity. We work with modules over a polynomial ring graded by a finitely generated abelian group. Imitating [14], our definition of regularity involves the vanishing of certain graded components of local cohomology. We establish the key properties of regularity: its connection with the minimal generators of a module and its behavior in short exact sequences. As a consequence, we are able relate the regularity of a module to chain complexes associated with the module.

Our primary motivation for studying regularity over multigraded polynomial rings comes from toric geometry. For a simplicial toric variety X , the homogeneous coordinate ring, introduced by Cox in [11], is a polynomial ring S graded by the divisor class group G of X . The dictionary linking the geometry of X with the theory of G -graded S -modules leads to geometric interpretations and applications for multigraded regularity. We prove that the regularity of an ideal sheaf bounds the multidegrees of the equations that cut out

the corresponding subvariety. Multigraded regularity also supplies a criterion for testing if an ample line bundle on X determines a projectively normal embedding.

Multigraded regularity consolidates a range of existing ideas. In the standard graded case, it reduces to Castelnuovo-Mumford regularity. If S has a nonstandard \mathbb{Z} -grading, then our definition is the version of regularity introduced by Benson in [3] for studying group cohomology. When S is the homogeneous coordinate ring of a product of projective spaces, multigraded regularity is the weak form of bigraded regularity defined by Hoffman and Wang in [25]. Our description for the multigraded regularity of fat points (Proposition 6.7) is also connected with the results in [21]. On the other hand, the versions of regularity developed for Grassmannians in [9] and abelian varieties in [33] do not fall under the umbrella of multigraded regularity.

For ease of exposition we state our theorems in the case where S is the homogeneous coordinate ring of a smooth projective toric variety X . Let B be the irrelevant ideal of X . We denote by $\mathbb{N}\mathcal{C}$ the semigroup generated by a finite subset $\mathcal{C} = \{c_1, \dots, c_\ell\}$ of $G \cong \text{Pic}(X)$. In the introduction, we restrict to the special case that \mathcal{C} is the minimal generating set of the semigroup of nef line bundles on X . For example, if $X = \mathbb{P}^d$ then $G \cong \mathbb{Z}$, S has the standard grading defined by $\deg(x_i) = \mathbf{1}$ for $1 \leq i \leq n$, $B = \langle x_1, \dots, x_n \rangle$, and $\mathcal{C} = \{\mathbf{1}\}$.

The main point of this paper is to introduce the following definition, which generalizes Castelnuovo-Mumford regularity.

Definition 1.1. For $m \in G$, we say that a G -graded S -module M is m -regular (with respect to \mathcal{C}) if the following conditions are satisfied:

- (1) $H_B^i(M)_p = 0$ for all $i \geq 1$ and all $p \in \bigcup (m - \lambda_1 c_1 - \dots - \lambda_\ell c_\ell + \mathbb{N}\mathcal{C})$ where the union is over all $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$ such that $\lambda_1 + \dots + \lambda_\ell = i - 1$.
- (2) $H_B^0(M)_p = 0$ for all $p \in \bigcup_{1 \leq j \leq \ell} (m + c_j + \mathbb{N}\mathcal{C})$.

We set $\text{reg}(M) := \{p \in G : M \text{ is } p\text{-regular}\}$.

In contrast with the usual notation, $\text{reg}(M)$ is a set rather than a single group element. Traditionally, $G = \mathbb{Z}$ and the regularity of M refers to the smallest $m \in G$ such that M is m -regular. When S has a multigrading, the group G is not equipped with a natural ordering so one cannot choose a smallest degree m . More significantly, the set $\text{reg}(M)$ may not even be determined by a single element; see Example 1.2. From this vantage point, bounding the regularity of a module M is equivalent to giving a subset of $\text{reg}(M)$.

Example 1.2. When X is the Hirzebruch surface \mathbb{F}_2 , the homogeneous coordinate ring $S = \mathbb{k}[x_1, x_2, x_3, x_4]$ has the \mathbb{Z}^2 -grading defined by $\deg(x_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\deg(x_2) = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$, $\deg(x_3) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\deg(x_4) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ and the irrelevant ideal B is $\langle x_1 x_2, x_2 x_3, x_3 x_4, x_1 x_4 \rangle$. This grading identifies $\text{Pic}(X)$ with \mathbb{Z}^2 . It follows that the semigroup of nef line bundles is generated by the set $\mathcal{C} = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$. A topological interpretation for $H_B^i(S)$ (see Section

3) shows that $\text{reg}(S) = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} + \mathbb{N}^2 \right) \cup \left(\begin{bmatrix} 0 \\ 1 \end{bmatrix} + \mathbb{N}^2 \right)$. A graphical representation of $\text{reg}(S)$ appears in Figure 5. Observe that $\text{reg}(S)$ is not determined by a single element of G and $\begin{bmatrix} 0 \\ 0 \end{bmatrix} \notin \text{reg}(S)$.

The following key result shows that the regularity of a module is a cohomological approximation for the degrees of its minimal generators.

Theorem 1.3. *The degrees of the minimal generators of a finitely generated G -graded S -module M do not belong to the set $\text{reg}(M) + \left(\bigcup_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathbb{N}\mathcal{C}) \right)$.*

To emphasize the similarities between the multigraded form of regularity and the original definition in [30], §14, we give the geometric translation of this result below. For $\mathbf{p} \in G$, $\mathcal{O}_X(\mathbf{p})$ is the associated line bundle on X . Set $\mathcal{F}(\mathbf{p}) := \mathcal{F} \otimes \mathcal{O}_X(\mathbf{p})$.

Theorem 1.4. *Let \mathcal{F} be a coherent \mathcal{O}_X -module. If \mathcal{F} is \mathbf{m} -regular then for every $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$ we have the following:*

- (1) \mathcal{F} is \mathbf{p} -regular.
- (2) The natural map $H^0(X, \mathcal{F}(\mathbf{p})) \otimes H^0(X, \mathcal{O}_X(\mathbf{q})) \rightarrow H^0(X, \mathcal{F}(\mathbf{p} + \mathbf{q}))$ is surjective for all $\mathbf{q} \in \mathbb{N}\mathcal{C}$.
- (3) $\mathcal{F}(\mathbf{p})$ is generated by its global sections.

We highlight two consequences of this theorem. Firstly, if \mathcal{I} is an ideal sheaf on X and $\mathbf{m} \in \text{reg}(\mathcal{I})$ then the subscheme of X defined by \mathcal{I} is cut out by equations of degree \mathbf{m} . Secondly, if $\mathcal{O}_X(\mathbf{p})$ is an ample line bundle and $\mathbf{p} \in \text{reg}(\mathcal{O}_X)$ then the complete linear series associated to $\mathcal{O}_X(\mathbf{p})$ gives a projectively normal embedding of X . In particular, if $\mathbf{0} \in \text{reg}(\mathcal{O}_X)$ then every ample line bundle on X gives a projectively normal embedding.

The next result illustrates a second key feature of regularity, namely its behavior in exact sequences. When S has the standard grading, the following are equivalent: the module M is \mathbf{m} -regular, the degrees of the i th syzgies are at most $\mathbf{m} + i$, and the truncated module $M|_{\geq \mathbf{m}}$ has a linear resolution. We generalize these properties in the following way.

Theorem 1.5. *Let d be the number of variable in polynomial ring S minus the rank of group G and let M be a finitely generated G -graded S -module.*

(1) *If $\cdots \rightarrow E_3 \rightarrow E_2 \rightarrow E_1 \rightarrow E_0 \xrightarrow{\hat{\partial}_0} M \rightarrow 0$ is a chain complex of finitely generated G -graded S -modules with B -torsion homology and $\hat{\partial}_0$ is surjective then*

$$\bigcup_{\phi: [d+1] \rightarrow [\ell]} \left(\bigcap_{1 \leq i \leq d+1} (-\mathbf{c}_{\phi(1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}(E_i)) \right) \subseteq \text{reg}(M).$$

(2) *If $\mathbf{c} \in \text{reg}(S) \cap \left(\bigcap_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathbb{N}\mathcal{C}) \right)$ and $\mathbf{m} \in \text{reg}(M)$ then there exists a chain complex*

$$\cdots \rightarrow E_3 \rightarrow E_2 \rightarrow E_1 \rightarrow E_0 \rightarrow M \rightarrow 0$$

with B -torsion homology and $E_i = \bigoplus S(-\mathbf{m} - ic)$.

If S has the standard grading, then the inclusion in Part (1) is an equality when the E_i form a minimal free resolution of M , and the chain complex in Part (2) is the minimal free resolution of $M|_{\geq m}$. Since B -torsion modules correspond to the zero sheaf on the corresponding toric variety, there is also a geometric version of this theorem involving regularity and resolutions of a sheaf.

The techniques used to establish Theorems 1.3, 1.4 and 1.5 also apply to a larger class of multigraded polynomial rings. We develop these methods for pairs (S, B) where S is a polynomial ring graded by a finitely generated abelian group G , and B is a monomial ideal that encodes a certain combinatorial structure of the grading. This class of rings includes the homogeneous coordinate rings of simplicial semi-projective toric varieties. We also establish these results for other choices of the set \mathcal{C} .

This paper is organized as follows. In the next section, we discuss the basic definitions and examples of multigraded polynomial rings. In Section 3, we establish some vanishing theorems for local cohomology modules. These results are based on a topological description for the multigraded Hilbert series of $H_B^i(S)$. The definition of multigraded regularity is presented in Section 4. We also prove that in certain cases the definition of regularity is equivalent to an apparently weaker vanishing condition. Section 5 connects the multigraded regularity of a module with the degrees of its minimal generators. In Section 6, we reinterpret regularity of S -module in terms of coherent sheaves on a simplicial toric variety and study some geometric applications. Finally, Section 7 examines the relationship between chain complexes associated to a module or sheaf and regularity.

Acknowledgements. We thank Dave Benson for conversations which expanded our notion of regularity. We are also grateful to Kristina Crona for conversations about multigradings early in this project. The computer software package Macaulay 2 [19] was indispensable for computing examples. Both authors were partially supported by the Mathematical Sciences Research Institute in Berkeley, CA.

2. Multigraded polynomial rings

In this section, we develop the foundations of multigraded polynomial rings. Let \mathbb{k} be a field and let G be a finitely generated abelian group. Throughout this paper, we work with a pair (S, B) where $S := \mathbb{k}[x_1, \dots, x_n]$ is a G -graded polynomial ring and B is a monomial ideal in S . For a positive integer m , we write $[m]$ for the set $\{1, \dots, m\}$. The convex cone generated by (or positive hull of) the vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ is the set

$$\text{pos}(\mathbf{v}_1, \dots, \mathbf{v}_m) := \{\lambda_1 \mathbf{v}_1 + \cdots + \lambda_m \mathbf{v}_m : \lambda_1, \dots, \lambda_m \in \mathbb{R}_{\geq 0}\}.$$

A G -grading of the polynomial ring $S = \mathbb{k}[x_1, \dots, x_n]$ corresponds to a semigroup homomorphism $\mathbb{N}^n \rightarrow G$. We say a monomial $\mathbf{x}^{\mathbf{u}}$ has degree \mathbf{p} if $\mathbf{u} \mapsto \mathbf{p} \in G$. This map induces a decomposition $S = \bigoplus_{\mathbf{p} \in G} S_{\mathbf{p}}$ satisfying $S_{\mathbf{p}} \cdot S_{\mathbf{q}} \subseteq S_{\mathbf{p}+\mathbf{q}}$ where $S_{\mathbf{p}}$ is the \mathbb{k} -span of

all $\mathbf{x}^{\mathbf{a}}$ of degree \mathbf{p} . The G -grading is determined by the set $\mathcal{A} := \{\mathbf{a}_1, \dots, \mathbf{a}_n\}$ where $\mathbf{a}_i := \deg(x_i) \in G$ for $1 \leq i \leq n$. We write $\mathbb{N}\mathcal{A}$ for the subsemigroup of G generated by \mathcal{A} . Let r be the rank of G and identify $\mathbb{R} \otimes_{\mathbb{Z}} G$ with \mathbb{R}^r . If $\bar{\mathbf{a}}_i$ denotes the image of \mathbf{a}_i in $\mathbb{R} \otimes_{\mathbb{Z}} G = \mathbb{R}^r$, then the set $\bar{\mathcal{A}} := \{\bar{\mathbf{a}}_1, \dots, \bar{\mathbf{a}}_n\}$ is an integral vector configuration in \mathbb{R}^r . The monomial ideal B corresponds to a chamber (maximal cell) $\Gamma \subset \mathbb{R}^r$ in the chamber complex of the vector configuration $\bar{\mathcal{A}}$. The chamber complex is the coarsest fan with support $\text{pos}(\bar{\mathcal{A}})$ that refines all the triangulations of $\bar{\mathcal{A}}$; see [4]. We encode the choice of a chamber Γ in the monomial ideal

$$B := \left\langle \prod_{i \in \sigma} x_i : \sigma \subseteq [n] \text{ with } \Gamma \subseteq \text{pos}(\bar{\mathbf{a}}_i : i \in \sigma) \right\rangle.$$

Alternatively, the monomial ideal B can be described by a regular triangulation ([36], Definition 5.3) of the dual vector configuration. If $d := n - r$, then the dual vector configuration is a set of vectors $\bar{\mathcal{B}} := \{\bar{\mathbf{b}}_1, \dots, \bar{\mathbf{b}}_n\}$ in \mathbb{R}^d such that

$$(2.0.1) \quad 0 \longrightarrow \mathbb{R}^d \xrightarrow{[\bar{\mathbf{b}}_1 \cdots \bar{\mathbf{b}}_n]^T} \mathbb{R}^n \xrightarrow{[\bar{\mathbf{a}}_1 \cdots \bar{\mathbf{a}}_n]} \mathbb{R}^r \longrightarrow 0$$

is a short exact sequence. The set $\bar{\mathcal{B}}$ is uniquely determined up to a linear change of coordinates on \mathbb{R}^d ; see [36], §6.4. We identify a triangulation of $\bar{\mathcal{B}}$ with a simplicial complex Δ . Gale duality, specifically [4], Theorem 3.1, implies that the chamber $\Gamma \subset \mathbb{R}^r$ corresponds to a regular triangulation Δ of $\bar{\mathcal{B}}$. For $\sigma \subseteq [n]$, let $\hat{\sigma}$ denote the complement of σ in $[n]$. From this standpoint, we have $B = \left\langle \prod_{i \in \hat{\sigma}} x_i : \sigma \in \Delta \right\rangle$.

The simplicial complex Δ (or equivalently the chamber Γ) also gives rise to two important subsemigroups of $\mathbb{N}\mathcal{A}$. The first subsemigroup \mathcal{K} is the intersection $\bigcap_{\sigma \in \Delta} \mathbb{N}\mathcal{A}_{\hat{\sigma}}$ where $\mathcal{A}_{\hat{\sigma}} := \{\mathbf{a}_i : i \in \hat{\sigma}\}$. Let $\mathbb{Z}\mathcal{A}$ be the subgroup of G generated by \mathcal{A} . The second subsemigroup \mathcal{K}^{sat} is the saturation (or normalization) of \mathcal{K} in $\mathbb{Z}\mathcal{A}$. In other words, \mathcal{K}^{sat} is the set of all $\mathbf{p} \in \mathbb{N}\mathcal{A}$ such that the image $\bar{\mathbf{p}} \in \mathbb{R}^r$ lies in the chamber Γ . The interior of \mathcal{K}^{sat} , denoted $\text{int } \mathcal{K}^{\text{sat}}$, consists of all $\mathbf{p} \in \mathcal{K}^{\text{sat}}$ such that $\bar{\mathbf{p}}$ lies in the interior of Γ .

Example 2.1. Let $G = \mathbb{Z}$ and assume that $\mathbf{a}_i > 0$ for all i . Since G is torsion-free, we may identify \mathbf{a}_i with $\bar{\mathbf{a}}_i$. The polynomial ring $S = \mathbb{k}[x_1, \dots, x_n]$ has the \mathbb{Z} -grading induced by $\deg(x_i) = \mathbf{a}_i$ for $1 \leq i \leq n$. The chamber complex of \mathcal{A} has a unique maximal cell $\Gamma = \mathbb{R}_{\geq 0}$ and the corresponding ideal is $B = \langle x_1, \dots, x_n \rangle$. The dual vector configuration $\bar{\mathcal{B}}$ is given by the columns of the matrix

$$\begin{bmatrix} \mathbf{a}_n & 0 & \cdots & 0 & -\mathbf{a}_1 \\ 0 & \mathbf{a}_n & \cdots & 0 & -\mathbf{a}_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \mathbf{a}_n & -\mathbf{a}_{n-1} \end{bmatrix}.$$

The semigroup $\mathcal{K} \subset \mathbb{Z}$ consists of all nonnegative multiples of $\text{lcm}(\mathcal{A})$ and $\mathcal{K}^{\text{sat}} = \mathbb{N}$. It follows that $\mathcal{K}^{\text{sat}} = \mathcal{K}$ if and only if $\mathbf{a}_i = 1$ for all $1 \leq i \leq n$. When $\mathbf{a}_i = 1$ for all i , S is standard graded polynomial ring and B is the unique graded maximal ideal.

Example 2.2. Fix $t \in \mathbb{N}$. Suppose that $G = \mathbb{Z}^2$ and let \mathcal{A} correspond to the columns of the matrix

$$\begin{bmatrix} 1 & -t & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}.$$

Again, we identify \mathcal{A} with $\bar{\mathcal{A}}$. The polynomial ring $S = \mathbb{k}[x_1, x_2, x_3, x_4]$ has the \mathbb{Z}^2 -grading defined by $\deg(x_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\deg(x_2) = \begin{bmatrix} -t \\ 0 \end{bmatrix}$, $\deg(x_3) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\deg(x_4) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. There are two possible choices for the chamber $\Gamma: \mathbb{R}_{\geq 0}^2 = \text{pos}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$ or $\text{pos}\left(\begin{bmatrix} -t \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$ and the ideal B equals $\langle x_1x_2, x_2x_3, x_3x_4, x_1x_4 \rangle$ or $\langle x_1x_2, x_2x_3, x_2x_4 \rangle$ respectively. The dual vector configuration is given by the columns of the matrix

$$\begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & t & -1 \end{bmatrix}.$$

Figure 1 illustrates the associated vector configurations when $t = 2$. Regardless of the choice of Γ , both \mathcal{H}^{sat} and \mathcal{H} equal $\Gamma \cap \mathbb{Z}^2$.




Figure 1. The vector configurations for Example 2.2.

Example 2.3. If $G = \mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^2$ and \mathcal{A} corresponds to the columns of the matrix

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \end{bmatrix}$$

where the entries in bottom two rows are elements of $\mathbb{Z}/2\mathbb{Z}$, then $S = \mathbb{k}[x_1, x_2, x_3, x_4, x_5]$ is graded by a group with torsion. The vector configuration $\bar{\mathcal{A}}$ is given by the columns of the matrix $\begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 \end{bmatrix}$. The chamber Γ is either $\Gamma_1 := \text{pos}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}\right)$ or $\Gamma_2 := \text{pos}\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}\right)$ and the ideal B equals either $\langle x_1x_2, x_1x_4, x_1x_5, x_2x_3, x_3x_4, x_3x_5 \rangle$ in the first case or $\langle x_1x_2, x_2x_3, x_2x_5, x_1x_4, x_3x_4, x_4x_5 \rangle$ in the second. The dual vector configuration $\bar{\mathcal{B}}$ is given by the columns of the matrix

$$\begin{bmatrix} 1 & 1 & -1 & -1 & 0 \\ 1 & -1 & -1 & 1 & 0 \\ 1 & 1 & 1 & 1 & -2 \end{bmatrix}.$$

The two triangulations are illustrated in Figure 2. If Γ_1 is the chosen chamber, then

$$\mathcal{K} = \mathbb{N} \left\{ \begin{bmatrix} 2 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 0 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{K}^{\text{sat}} = \mathbb{N} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$




Figure 2. The two triangulations and chamber complex for Example 2.3.

On the other hand, if Γ_2 is the chosen chamber, then

$$\mathcal{K} = \mathbb{N} \left\{ \begin{bmatrix} 0 \\ 2 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 0 \\ 0 \end{bmatrix} \right\} \quad \text{and} \quad \mathcal{K}^{\text{sat}} = \mathbb{N} \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

The following lemma further illustrates the connections between the ideal B and the semigroups \mathcal{K} and \mathcal{K}^{sat} . For $\mathbf{p} \in G$, let $\langle S_{\mathbf{p}} \rangle$ denote the ideal generated by all the monomials of degree \mathbf{p} in S . For $\mathbf{u} \in \mathbb{N}^n$, let $\text{supp}(\mathbf{u}) := \{i : u_i \neq 0\} \subseteq [n]$.

Lemma 2.4. *If \mathbf{p} belongs to the interior of \mathcal{K}^{sat} and $\deg(\mathbf{x}^{\mathbf{u}}) = \mathbf{p}$ then $\mathbf{x}^{\mathbf{u}}$ belongs to the ideal B . Moreover, if $\mathbf{p} \in \mathcal{K}$, then $B \subseteq \sqrt{\langle S_{\mathbf{p}} \rangle}$.*

Proof. If $\mathbf{p} \in \text{int } \mathcal{K}^{\text{sat}}$ and $\mathbf{x}^{\mathbf{u}} \in S_{\mathbf{p}}$ then $\text{pos}(\bar{\mathbf{a}}_i : i \in \text{supp}(\mathbf{u})) \cap \text{int } \Gamma \neq \emptyset$. Since Γ is a maximal cell in the chamber complex of \mathcal{A} , we have $\Gamma \subseteq \text{pos}(\bar{\mathbf{a}}_i : i \in \text{supp}(\mathbf{u}))$ and $\dim_{\mathbb{R}} \text{pos}(\bar{\mathbf{a}}_i : i \in \text{supp}(\mathbf{u})) = r$. Caratheodory’s Theorem ([36], Proposition 1.15) implies that $\text{pos}(\bar{\mathbf{a}}_i : i \in \text{supp}(\mathbf{u}))$ is the union of the $\text{pos}(\bar{\mathbf{a}}_i : i \in \sigma)$ where $|\sigma| = r$ and $\sigma \subseteq \text{supp}(\mathbf{u})$. Hence, there exists $\sigma \subseteq \text{supp}(\mathbf{u})$ such that $|\sigma| = r$ and $\text{pos}(\bar{\mathbf{a}}_i : i \in \sigma) \cap \text{int } \Gamma \neq \emptyset$. Again, because Γ is a chamber, we have $\Gamma \subseteq \text{pos}(\bar{\mathbf{a}}_i : i \in \sigma)$. It follows that the monomial $\prod_{i \in \sigma} x_i$ belongs to B and divides $\mathbf{x}^{\mathbf{u}}$. This establishes the first assertion.

If $\mathbf{p} \in \mathcal{H}$, then for every $\sigma \in \Delta$ there exists a monomial $\mathbf{x}^{\mathbf{u}} \in S_{\mathbf{p}}$ with $\text{supp}(\mathbf{u}) \subseteq \hat{\sigma}$. It follows that a sufficiently large power of each generator of B belongs to the ideal $\langle S_{\mathbf{p}} \rangle$ which implies $B \subseteq \sqrt{\langle S_{\mathbf{p}} \rangle}$. \square

Our motivating example of a pair (S, B) is the homogeneous coordinate ring and irrelevant ideal of a toric variety introduced by Cox in [11]. Let X be a simplicial toric variety over a field \mathbb{k} determined by a fan Δ in \mathbb{R}^d . By numbering the rays (one-dimensional cones) of Δ , we identify Δ with a simplicial complex on $[n]$. We write $\mathbf{b}_1, \dots, \mathbf{b}_n$ for the unique minimal lattice vectors generating the rays and we assume that the positive hull of $\mathcal{B} := \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$ is \mathbb{R}^d . The set \mathcal{B} gives rise to a short exact sequence

$$0 \longrightarrow \mathbb{Z}^d \xrightarrow{[\mathbf{b}_1 \cdots \mathbf{b}_n]^T} \mathbb{Z}^n \longrightarrow G \longrightarrow 0.$$

Tensoring this sequence with \mathbb{R} , we obtain (2.0.1). Geometrically, G is the divisor class group (or Chow group) of X ; see [17], §3.4. The image of the i th standard basis vector of \mathbb{Z}^n in G is denoted by \mathbf{a}_i . Observe that the set $\mathcal{A} = \{\mathbf{a}_1, \dots, \mathbf{a}_n\}$ is uniquely determined up to an automorphism of G . The *homogeneous coordinate ring* of X is the polynomial ring $S = \mathbb{k}[x_1, \dots, x_n]$ with the G -grading induced by $\deg(x_i) = \mathbf{a}_i$ and the *irrelevant ideal* is $B = \left\langle \prod_{i \notin \sigma} x_i : \sigma \in \Delta \right\rangle$.

This geometric choice of a pair (S, B) fits into the algebraic framework developed at the start of this section if the fan of X corresponds to a regular triangulation of \mathcal{B} . By [24], Theorem 2.6, this is equivalent to saying that X is semi-projective. In particular, this holds whenever X is projective. Conversely, an algebraic pair (S, B) is the homogeneous coordinate ring and irrelevant ideal of a simplicial toric variety if and only if, for each i , the ray $\text{pos}(\bar{\mathbf{b}}_i)$ belongs to the triangulation associated to the chamber Γ .

Example 2.5. The pair (S, B) described in Example 2.1 corresponds to the weighted projective space $X = \mathbb{P}(\mathbf{a}_1, \dots, \mathbf{a}_n)$. In particular, when $X = \mathbb{P}^d$, the homogeneous coordinate ring S is the standard graded polynomial ring and the irrelevant ideal B is the unique graded maximal ideal.

Example 2.6. If (S, B) is the pair described in Example 2.2 where $\Gamma = \mathbb{R}_{\geq 0}^2$ then the associated toric variety is the Hirzebruch surface (or rational scroll) $\mathbb{F}_t = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(t))$.

Example 2.7. If X is the product of projective space $\mathbb{P}^d \times \mathbb{P}^e$, then the homogeneous coordinate ring $S = \mathbb{k}[x_0, \dots, x_d, y_0, \dots, y_e]$ has the \mathbb{Z}^2 -grading induced by $\deg(x_i) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\deg(y_i) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ and the irrelevant ideal B is $\langle x_0, \dots, x_d \rangle \cap \langle y_0, \dots, y_e \rangle$.

When the pair (S, B) corresponds to a simplicial toric variety X , the chamber Γ and the semigroups \mathcal{H} and \mathcal{H}^{sat} have geometric interpretations. Assuming all the maximal cones are d -dimensional, we have $\text{Pic}(X) \otimes \mathbb{Q} \cong G \otimes \mathbb{Q}$. As [12], §3 indicates, the chamber Γ is the closure of the Kähler cone of X . The dual of the Kähler cone is the Mori cone of effective 1-cycles modulo numerical equivalence. The semigroup \mathcal{H}^{sat} corresponds to the numerically effective Weil divisors on X up to rational equivalence and elements in \mathcal{H} are nef line bundles. When X is smooth, the class group G is torsion-free and $\mathcal{H} = \mathcal{H}^{\text{sat}}$.

The next example demonstrates that \mathcal{H} can be complicated even when X is smooth.

Example 2.8. Consider the following smooth resolution X of weighted projective space $\mathbb{P}(2, 3, 7, 1)$. Specifically, the set \mathcal{B} corresponds to the columns of the matrix

$$\begin{bmatrix} 1 & 0 & 0 & -2 & 0 & 0 & -1 & 0 & -1 & -1 & -1 \\ 0 & 1 & 0 & -3 & 0 & -1 & -1 & -1 & -1 & -2 & -2 \\ 0 & 0 & 1 & -7 & -1 & -2 & -3 & -3 & -4 & -4 & -5 \end{bmatrix}.$$

Figure 3 illustrates the regular triangulation and the irrelevant ideal is

$$\begin{aligned} B = \langle & x_1x_2x_3x_4x_6x_7x_9x_{10}, & x_2x_3x_4x_5x_6x_7x_9x_{10}, & x_1x_2x_3x_6x_7x_8x_9x_{10}, \\ & x_2x_3x_5x_6x_7x_8x_9x_{10}, & x_1x_2x_3x_5x_7x_8x_9x_{11}, & x_1x_2x_4x_5x_7x_8x_9x_{11}, \\ & x_1x_2x_5x_6x_7x_8x_9x_{11}, & x_1x_2x_3x_6x_7x_8x_{10}x_{11}, & x_1x_3x_4x_6x_7x_8x_{10}x_{11}, \\ & x_1x_3x_5x_6x_7x_8x_{10}x_{11}, & x_2x_3x_4x_6x_7x_9x_{10}x_{11}, & x_1x_2x_5x_6x_8x_9x_{10}x_{11}, \\ & x_1x_3x_5x_6x_8x_9x_{10}x_{11}, & x_1x_4x_5x_6x_8x_9x_{10}x_{11}, & x_2x_3x_5x_7x_8x_9x_{10}x_{11}, \\ & x_2x_4x_5x_7x_8x_9x_{10}x_{11}, & x_3x_4x_6x_7x_8x_9x_{10}x_{11}, & x_4x_5x_6x_7x_8x_9x_{10}x_{11} \rangle. \end{aligned}$$

In this case, Γ is an 8-dimensional cone with 16 extremal rays. Using Normaliz [8], we determine that the set of minimal generators for \mathcal{H} has 25 elements all of which lie on the boundary of Γ .




Figure 3. The triangulation for Example 2.8.

3. A topological formula for local cohomology

Throughout this paper, M denotes a finitely generated G -graded S -module. In this section we derive a vanishing theorem for the local cohomology modules $H_B^i(M)$ from a topological formula for $H_B^i(S)$. We refer to [5] for background information on local cohomology. A module M is B -torsion if $M = H_B^0(M)$. If M is B -torsion then $H_B^i(M) = 0$ for $i > 0$. For an element $g \in S$, we set

$$(0 :_M g) := \{f \in M : gf = 0\} = \text{Ker}(M \xrightarrow{g} M).$$

This submodule is zero when g is a nonzerodivisor on M . We say an element $g \in S$ is *almost a nonzerodivisor on M* if $(0 :_M g)$ is a B -torsion module.

Proposition 3.1. *Let \mathbb{k} be an infinite field. If $\mathbf{p} \in \mathcal{H}$ and $g \in S$ is a sufficiently general form of degree \mathbf{p} , then g is almost a nonzerodivisor on M . In other words, there is an open dense set of degree \mathbf{p} forms g for which $(0 :_M g)$ is B -torsion.*

Proof. The module $M' := (0 :_M g)$ is B -torsion if each element in M' is annihilated by some power of B . This is equivalent to saying that the localization $M'_P = 0$ for all prime ideals P in S not containing B . In other words, g is a nonzerodivisor on the localization M_P . Hence, it suffices to show that g is not contained in any of the associated primes of M except possibly those which contain B .

To accomplish this, observe that each prime ideal P in S which does not contain B intersects $S_{\mathbf{p}}$ in a proper subspace. Otherwise $\langle S_{\mathbf{p}} \rangle \subseteq P$ and Lemma 2.4 implies that $B \subseteq \sqrt{\langle S_{\mathbf{p}} \rangle} \subseteq \sqrt{P} = P$ which contradicts the hypothesis on P . Because M has only finitely many associated primes, our observation shows that $g \in S_{\mathbf{p}}$ is almost a nonzerodivisor on M if it lies outside a certain finite union of proper subspaces. Since \mathbb{k} is infinite, the vector space $S_{\mathbf{p}}$ is not a finite union of proper subspaces. \square

To give a uniform vanishing result for local cohomology, we assume for the remainder of this section that $\text{pos}(\mathcal{A})$ is a pointed cone with $\bar{\mathbf{a}}_i \neq \mathbf{0}$ for each i or equivalently that \mathcal{A} is an acyclic vector configuration; see [36], §6.2. This condition holds if S is the homogeneous coordinate ring of a projective toric variety. We can rephrase this assumption by saying that the dual configuration $\bar{\mathcal{B}}$ is totally cyclic or $\text{pos}(\bar{\mathcal{B}}) = \mathbb{R}^d$. Hence, any regular triangulation of $\bar{\mathcal{B}}$ is a complete fan in \mathbb{R}^d . We may identify Δ with a simplicial $(d - 1)$ -sphere and the collection $\hat{\Delta} := \Delta \cup \{[n]\}$ (ordered by inclusion) with the face poset of a finite regular cell decomposition of a d -ball. It follows that there is an incidence function ε on $\hat{\Delta}$. Recall from [7], §6.2, that ε takes values in $\{0, \pm 1\}$ and is induced by an orientation of the cells.

For $\sigma \subseteq [n]$, let x_{σ} denote the squarefree monomial $\prod_{i \in \sigma} x_i \in S$. The *canonical Čech complex* associated to $\hat{\Delta}$ is the following chain complex:

$$\check{C} := 0 \longrightarrow \check{C}^0 \xrightarrow{\partial^0} \check{C}^1 \xrightarrow{\partial^1} \check{C}^2 \xrightarrow{\partial^2} \dots \xrightarrow{\partial_{d-1}} \check{C}^d \xrightarrow{\partial_d} \check{C}^{d+1} \longrightarrow 0,$$

where $\check{C}^0 := S$, $\check{C}^i := \bigoplus_{\sigma \in \Delta, |\sigma|=d+1-i} S[\mathbf{x}_{\sigma}^{-1}]$ for $i > 0$ and $\partial^i : \check{C}^i \rightarrow \check{C}^{i+1}$ is composed of homomorphisms $\varepsilon(\sigma, \tau) \cdot \text{nat} : S[\mathbf{x}_{\sigma}^{-1}] \rightarrow S[\mathbf{x}_{\tau}^{-1}]$. By combining [2], Corollary 2.13, and [29], Theorem 6.2, we deduce that $H_B^i(M) = H^i(\check{C} \otimes_S M)$ for all S -modules M . It follows that $H_B^i(M) = 0$ for all $i < 0$ or $i > d + 1$.

The next proposition gives a combinatorial description for the local cohomology of the polynomial ring S . We will use this formula to compute the multigraded regularity in some examples. This result is a variant on the formulae found in [16] and [34]. For $\sigma \in \Delta$, let Δ_{σ} be the induced subcomplex $\{\tau \in \Delta : \tau \subseteq \sigma\}$ of Δ . The modules $H_B^i(S)$ have a \mathbb{Z}^n -grading which refines their G -grading.

Proposition 3.2. *Let $\tilde{H}^i(-)$ denote the i th reduced cellular cohomology group with coefficients in \mathbb{k} . If $\mathbf{u} \in \mathbb{Z}^n$ and $\sigma := \text{neg}(\mathbf{u}) = \{j : u_j < 0\}$, then we have*

$$H_B^i(S)_{\mathbf{u}} \cong \begin{cases} \tilde{H}^{i-2}(\Delta_\sigma) & \text{for } i \neq 1, \\ 0 & \text{for } i = 1. \end{cases}$$

Proof. One easily checks (see [7], Lemma 5.3.6) that $\dim_{\mathbb{k}} S[x_{\hat{\tau}}^{-1}]_{\mathbf{u}} = 1$ when $\tau \subseteq \hat{\sigma}$ and $\dim_{\mathbb{k}} S[x_{\hat{\tau}}^{-1}]_{\mathbf{u}} = 0$ when $\tau \not\subseteq \hat{\sigma}$. It follows that the \mathbf{u} th graded component of $\tilde{\mathbf{C}}$ is isomorphic to a shift of the augmented oriented chain complex of $\hat{\Delta}_{\hat{\sigma}}$ (see [7], §6.2). More precisely, we have $H^i(\tilde{\mathbf{C}}_{\mathbf{u}}) = \tilde{H}_{d-i}(\hat{\Delta}_{\hat{\sigma}})$. To complete the proof, we analyze three cases:

- If σ is a proper nonempty subset of $[n]$, then the subcomplex $\hat{\Delta}_{\hat{\sigma}}$ equals $\Delta_{\hat{\sigma}}$. Alexander duality ([31], Theorem 71.1) shows that $\tilde{H}_{d-i}(\hat{\Delta}_{\hat{\sigma}})$ is isomorphic to $\tilde{H}^{i-2}(\Delta \setminus \Delta_{\hat{\sigma}})$. Because Δ_{σ} is a deformation retract of $\Delta \setminus \Delta_{\hat{\sigma}}$ ([31], Lemma 70.1), we have

$$\tilde{H}_{d-i}(\hat{\Delta}_{\hat{\sigma}}) \cong \tilde{H}^{i-2}(\Delta_{\sigma}).$$

Note that $\tilde{H}^{i-2}(\Delta_{\sigma}) = 0$ when $i = 1$.

- If $\sigma = \emptyset$, then $\hat{\sigma} = [n]$ and $\hat{\Delta}_{\hat{\sigma}}$ is a d -ball which implies that $\tilde{H}_{d-i}(\hat{\Delta}_{\hat{\sigma}}) = 0$ for all i . It follows that $\tilde{H}_{d-i}(\hat{\Delta}_{[n]})$ is isomorphic to $\tilde{H}^{i-2}(\Delta_{\emptyset})$ for $i \neq 1$ and equals 0 for $i = 1$.

- If $\sigma = [n]$, then $\hat{\sigma} = \emptyset$ and $\hat{\Delta}_{\hat{\sigma}} = \emptyset$ which implies that $\tilde{H}_{d-i}(\hat{\Delta}_{\hat{\sigma}}) = 0$ for $i \neq d + 1$ and $\tilde{H}_{-1}(\hat{\Delta}_{\hat{\sigma}}) \cong \mathbb{k}$. Since $\Delta_{[n]} = \Delta$ is a $(d - 1)$ -sphere, we also have $\tilde{H}_{d-i}(\hat{\Delta}_{\emptyset}) \cong \tilde{H}^{i-2}(\Delta_{[n]})$ for all i . \square

Remark 3.3. Proposition 3.2 implies that $H_B^i(S) \neq 0$ only if $2 \leq i \leq d + 1$. Since $H_B^{d+1}(S)_{\mathbf{u}} = \mathbb{k}$ when $\text{neg}(\mathbf{u}) = [n]$, we have $H_B^{d+1}(S) \neq 0$. Moreover, $H_B^{d+1}(S)$ is the only nonvanishing local cohomology module if and only if every proper subcomplex of Δ is contractible. This happens precisely when Δ is the boundary of the standard simplex. Hence, $H_B^i(S) = 0$ for all $i \neq d + 1$ if and only if $B = \langle x_1, \dots, x_n \rangle$.

Using Proposition 3.2, we can describe the Hilbert series of the modules $H_B^i(S)$.

Corollary 3.4. *For all $i \neq 1$, we have*

$$(3.4.1) \quad \sum_{\mathbf{p} \in G} \dim_{\mathbb{k}} H_B^i(S)_{\mathbf{p}} \cdot \mathbf{t}^{\mathbf{p}} = \sum_{\sigma \subseteq [n]} \frac{\dim_{\mathbb{k}} \tilde{H}^{i-2}(\Delta_{\sigma}) \cdot \prod_{j \in \sigma} \mathbf{t}^{-a_j}}{\prod_{j \in \hat{\sigma}} (1 - \mathbf{t}^{a_j}) \prod_{j \in \sigma} (1 - \mathbf{t}^{-a_j})}.$$

Proof. For $i \neq 1$, Proposition 3.2 implies that

$$\begin{aligned} \sum_{\mathbf{u} \in \mathbb{Z}^n} \dim_{\mathbb{k}} H_B^i(S)_{\mathbf{u}} \cdot \mathbf{x}^{\mathbf{u}} &= \sum_{\mathbf{u} \in \mathbb{Z}^n} \dim_{\mathbb{k}} \tilde{H}^{i-2}(\Delta_{\text{neg}(\mathbf{u})}) \cdot \mathbf{x}^{\mathbf{u}} \\ &= \sum_{\sigma \subseteq [n]} \left(\dim_{\mathbb{k}} \tilde{H}^{i-2}(\Delta_{\sigma}) \sum_{\text{neg}(\mathbf{u}) = \sigma} \mathbf{x}^{\mathbf{u}} \right) = \sum_{\substack{\sigma \subseteq [n] \\ j \in \hat{\sigma}}} \frac{\dim_{\mathbb{k}} \tilde{H}^{i-2}(\Delta_{\sigma}) \cdot \prod_{j \in \sigma} x_j^{-1}}{\prod_{j \in [n]} (1 - x_j) \prod_{j \in \sigma} (1 - x_j^{-1})}. \end{aligned}$$

Mapping x_j to \mathbf{t}^{a_j} establishes the corollary. \square

We illustrate this corollary with the following example.

Example 3.5. If S is the homogeneous coordinate ring of the Hirzebruch surface \mathbb{F}_t as in Examples 2.2 and 2.6, then the only non-contractible subcomplexes are Δ_\emptyset , $\Delta_{\{1,3\}}$, $\Delta_{\{2,4\}}$ and $\Delta_{[4]}$. Since $\tilde{H}^i(\Delta_\emptyset) = 0$ for all $i \neq -1$ and $\tilde{H}^{-1}(\Delta_\emptyset) = \mathbb{k}$, this subcomplex does not contribute to any local cohomology module. Because $\tilde{H}^0(\Delta_{\{1,3\}}) = \mathbb{k} = \tilde{H}^0(\Delta_{\{2,4\}})$, the degrees $\mathbf{p} \in \mathbb{Z}^2$ for which $H_B^2(S)_\mathbf{p} \neq 0$ correspond to the lattice points in the two cones $-\mathbf{a}_1 - \mathbf{a}_3 + \text{pos}(-\mathbf{a}_1, \mathbf{a}_2, -\mathbf{a}_3, \mathbf{a}_4)$ and $-\mathbf{a}_2 - \mathbf{a}_4 + \text{pos}(\mathbf{a}_1, -\mathbf{a}_2, \mathbf{a}_3, -\mathbf{a}_4)$. We also have $\tilde{H}^1(\Delta_{[n]}) = \mathbb{k}$, so the degrees $\mathbf{p} \in \mathbb{Z}^2$ for which $H_B^2(S)_\mathbf{p} \neq 0$ correspond to the lattice points in the cone $-\mathbf{a}_1 - \mathbf{a}_2 - \mathbf{a}_3 - \mathbf{a}_4 + \text{pos}(-\mathbf{a}_1, -\mathbf{a}_2, -\mathbf{a}_3, -\mathbf{a}_4)$. These cones are indicated by the shaded areas in Figure 4 when $t = 2$.




Figure 4. Degrees $\mathbf{p} \in \mathbb{Z}^2$ for which $H_B^i(S)_\mathbf{p} \neq 0$ in Example 3.5.

The next result extends a well-known vanishing theorem for ample line bundles on a complete toric variety; see (6.3.1) for the explicit connection.

Corollary 3.6. *If \mathbf{p} belongs to \mathcal{K}^{sat} , then $H_B^i(S)_\mathbf{p}$ vanishes.*

Proof. Fix $\mathbf{p} \in \mathcal{K}^{\text{sat}}$ and suppose $\mathbf{u} \in \mathbb{Z}^n$ satisfies $\text{deg}(\mathbf{x}^{\mathbf{u}}) = \mathbf{p}$. The vector \mathbf{u} defines a function on $\psi_{\mathbf{u}} : \mathbb{R}^d \rightarrow \mathbb{R}$ that is linear on the cone $\text{pos}(\bar{\mathbf{b}}_i : i \in \sigma)$ for all $\sigma \in \Delta$ and satisfies $\psi_{\mathbf{u}}(\bar{\mathbf{b}}_i) = -u_i$ for $1 \leq i \leq n$. [4], Lemma 3.2 implies that $\mathbf{p} \in \mathcal{K}^{\text{sat}}$ if and only if $\psi_{\mathbf{u}}$ is convex which means $\psi_{\mathbf{u}}(\mathbf{w} + \mathbf{w}') \geq \psi_{\mathbf{u}}(\mathbf{w}) + \psi_{\mathbf{u}}(\mathbf{w}')$ for all $\mathbf{w}, \mathbf{w}' \in \mathbb{R}^d$. Hence, the set $\{\mathbf{w} \in \mathbb{R}^d : \psi_{\mathbf{u}}(\mathbf{w}) \geq 0\}$ is convex. It follows that simplicial complex $\Delta_{\text{neg}(\mathbf{u})}$ corresponds to the intersection of a convex set with the $(d - 1)$ -sphere. Therefore, $\Delta_{\text{neg}(\mathbf{u})}$ is contractible and Proposition 3.2 implies that $H_B^i(S)_\mathbf{p} = 0$ which proves the statement. \square

As a further corollary, we obtain a vanishing theorem for the local cohomology of any finitely generated G -graded S -module. Geometrically, this result corresponds to Fujita’s vanishing theorem ([27], Theorem 1.4.32). We first record a useful observation.

Remark 3.7. Let \mathcal{D} be a subsemigroup of G consisting of all points $\mathbf{p} \in G$ such that their image $\bar{\mathbf{p}}$ in \mathbb{R}^r lies in a fixed convex cone C . If $\mathbf{p} + \mathcal{D}$ and $\mathbf{q} + \mathcal{D}$ are two shifts of \mathcal{D} , then their intersection is nonempty. In fact, if $\mathbf{p} \in \mathcal{D}$, with $\bar{\mathbf{p}} \in \text{int } C$, then for all sufficiently large $j \in \mathbb{N}$ we have $j\mathbf{p} \in \mathbf{q} + \mathcal{D}$ for any $\mathbf{q} \in G$.

Corollary 3.8. *There is $\mathbf{m} \in \mathcal{K}^{\text{sat}}$ such that $H_B^i(M)_\mathbf{p} = 0$ for all i and all $\mathbf{p} \in \mathbf{m} + \mathcal{K}^{\text{sat}}$. In particular, $H_B^i(M)_\mathbf{p}$ vanishes for all i and all $\mathbf{p} \in \mathcal{K}^{\text{sat}}$ sufficiently far into the interior of \mathcal{K}^{sat} . If desired we may assume $\mathbf{m} \in \mathcal{K}$.*

Proof. We proceed by induction on the projective dimension $\text{pd}(M)$ of M . Since M is finitely generated, there is a short exact sequence $0 \rightarrow E_1 \rightarrow E_0 \rightarrow M \rightarrow 0$ where $E_0 = \bigoplus_{1 \leq j \leq h} S(-\mathbf{q}_j)$ for some $\mathbf{q}_j \in G$. The associated long exact sequence contains

$$(3.8.2) \quad H_B^i(E_0)_\mathbf{p} \rightarrow H_B^i(M)_\mathbf{p} \rightarrow H_B^{i+1}(E_1)_\mathbf{p}.$$

If we choose $\mathbf{m}' \in \bigcap_{1 \leq j \leq h} (\mathbf{q}_j + \mathcal{K}^{\text{sat}})$, which is possible by Remark 3.7, then Theorem 3.6 implies that the left module in (3.8.2) vanishes for all i and all $\mathbf{p} \in \mathbf{m}' + \mathcal{K}^{\text{sat}}$. When $\text{pd}(M) = 0$, we have $E_1 = 0$ and there is nothing more to prove. Otherwise, we have $E_1 \neq 0$ and $\text{pd}(E_1) < \text{pd}(M)$. Hence, the induction hypothesis provides $\mathbf{m}'' \in G$ such that the right module in (3.8.2) vanishes for all $\mathbf{p} \in \mathbf{m}'' + \mathcal{K}^{\text{sat}}$. Therefore, by choosing

$$\mathbf{m} \in (\mathbf{m}' + \mathcal{K}^{\text{sat}}) \cap (\mathbf{m}'' + \mathcal{K}^{\text{sat}}),$$

which is again possible by Remark 3.7, we see that the middle module in (3.8.2) vanishes for all i and all $\mathbf{p} \in \mathbf{m} + \mathcal{K}^{\text{sat}}$.

Lastly, because \mathcal{K}^{sat} is the saturation of \mathcal{K} , there exists some $\mathbf{m}''' \in \mathcal{K} \cap (\mathbf{m} + \mathcal{K}^{\text{sat}})$. Since $\mathbf{m}''' + \mathcal{K}^{\text{sat}} \subseteq \mathbf{m} + \mathcal{K}^{\text{sat}}$, the corresponding vanishing statement holds for \mathbf{m}''' which establishes the last part of the corollary. \square

We end this section by examining the assumption that $\text{pos}(\overline{\mathcal{A}})$ is acyclic. Firstly, we can remove the condition that $\overline{\mathbf{a}}_i \neq \mathbf{0}$ for all i .

Remark 3.9. Let $\sigma := \{i \in [n] : \overline{\mathbf{a}}_i = \mathbf{0}\}$. For each $j \in \sigma$ the variable x_j does not divide any minimal generator of B . In other words, the set σ is contained in every facet of Δ . Hence, the simplicial complex Δ is a cone over the induced subcomplex Δ_σ . By replacing Δ with Δ_σ , we can extend Proposition 3.2 and its corollaries to this more general situation.

On the other hand, the next example shows that we cannot eliminate the hypothesis that $\text{pos}(\overline{\mathcal{A}})$ is a pointed cone.

Example 3.10. If $G = \mathbb{Z}$ and $\mathcal{A} = \{\mathbf{1}, -\mathbf{1}, \mathbf{1}\}$, then $S = \mathbb{k}[x_1, x_2, x_3]$ has the \mathbb{Z} -grading induced by $\deg(x_2) = -\mathbf{1}$ and $\deg(x_1) = \deg(x_3) = \mathbf{1}$. Choosing the chamber Γ to be $\mathbb{R}_{\geq 0}$, we have $\mathcal{K} = \mathcal{K}^{\text{sat}} = \mathbb{N}$. The corresponding monomial ideal is $B = \langle x_1, x_3 \rangle$ and the dual vector configuration corresponds to the columns of the matrix $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$. Since the first paragraph of the proof of Proposition 3.2 applies in this situation,

$$H_B^2(S)_\mathbf{u} = \tilde{H}_0(\hat{\Delta}_\sigma) \quad \text{where } \sigma = \text{neg}(\mathbf{u}).$$

If $\sigma = \{2\}$, then $\hat{\sigma} = \{1, 3\}$ and $\hat{\Delta}_{\{1,3\}} = \{\emptyset, \{1\}, \{3\}\}$. It follows that $\tilde{H}_0(\hat{\Delta}_\sigma) \neq 0$ which implies that $H_B^2(S)_\mathbf{u} \neq 0$ when $\text{neg}(\mathbf{u}) = \{2\}$. However, for all $\mathbf{p} \in \mathbb{N}$, there exists a $\mathbf{u} \in \mathbb{Z}^3$ with $\text{neg}(\mathbf{u}) = \{2\}$. We conclude that $H_B^2(S)_\mathbf{p} \neq 0$ for all $\mathbf{p} \in \mathcal{K}$.

4. Regularity of S -modules

In this section, we define the multigraded regularity of a G -graded S -module. As in the standard case, multigraded regularity involves the vanishing of certain graded components of local cohomology modules. We show that, in certain cases, the definition is equivalent to an apparently weaker vanishing statement.

Before giving the definition of multigraded regularity, we collect some notation. Throughout this paper, $\mathcal{C} := \{c_1, \dots, c_\ell\}$ will be a fixed subset of G . We write $\mathbb{N}\mathcal{C}$ for the subsemigroup of G generated by \mathcal{C} . A subset \mathcal{D} of G is called an $\mathbb{N}\mathcal{C}$ -module if for all $p \in \mathcal{D}$ and all $q \in \mathbb{N}\mathcal{C}$ we have $p + q \in \mathcal{D}$. If $\mathcal{D} \subseteq G$ and $i \in \mathbb{Z}$, then

$$\mathcal{D}[i] := \bigcup \left(\frac{i}{|i|} (\lambda_1 c_1 + \dots + \lambda_\ell c_\ell) + \mathcal{D} \right) \subseteq G$$

where the union is over all $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$ such that $\lambda_1 + \dots + \lambda_\ell = |i|$. For $p \in G$, we clearly have $p + \mathcal{D}[i] = (p + \mathcal{D})[i]$. Moreover, if \mathcal{D} is an $\mathbb{N}\mathcal{C}$ -module then $\mathcal{D}[i]$ is also an $\mathbb{N}\mathcal{C}$ -module and $\mathcal{D}[i + 1] \subseteq \mathcal{D}[i]$. The following definition includes Definition 1.1 as a special case.

Definition 4.1. If $m \in G$, then the S -module M is *m -regular* (with respect to \mathcal{C}) if $H_B^i(M)_p = 0$ for all $i \geq 0$ and all $p \in m + \mathbb{N}\mathcal{C}[1 - i]$. The *regularity of M* , denoted $\text{reg}(M)$, is the set $\{m \in G : M \text{ is } m\text{-regular}\}$.

When S has the standard grading, $B = \langle x_1, \dots, x_n \rangle$, and $\mathcal{C} = \{\mathbf{1}\}$, the definition is simply: M is *m -regular* if and only if $H_B^i(M)_p = 0$ for all $i \geq 0$ and all $p \geq m - i + 1$. This is equivalent to the standard definition of Castelnuovo-Mumford regularity; for example see [5], page 282. Moreover, if S has an \mathbb{N} -grading as in Example 2.1, $B = \langle x_1, \dots, x_n \rangle$ and $\mathcal{C} = \{\mathbf{1}\}$, then the definition of *m -regular* is compatible with [3], Definition 4.1.

Our definition for $\text{reg}(M)$ conflicts with the standard notation used for Castelnuovo-Mumford regularity. Traditionally, $G = \mathbb{Z}$ and the regularity of M referred to the smallest $m \in G$ such that M is *m -regular*. In the multigraded setting, the group G is not equipped with a natural ordering so one cannot choose a smallest degree m . More importantly, the set $\text{reg}(M)$ may not even be determined by a single element of G . Example 4.3 illustrates this phenomenon. For these reasons, we regard the regularity of M as a subset of G . From this vantage point, giving a bound on the smallest m such that M is *m -regular* should be interpreted as describing a subset of $\text{reg}(M)$.

Example 4.2. Let $S = \mathbb{k}[x_1, \dots, x_n]$ have the \mathbb{Z} -grading induced by $\deg(x_i) = a_i > 0$ and the ideal B be $\langle x_1, \dots, x_n \rangle$ as in Example 2.1. If $\mathcal{C} = \{\mathbf{1}\}$, then Remark 3.3 implies that $\text{reg}(S) = \{m \in \mathbb{Z} : m \geq n - a_1 - \dots - a_n\}$. On the other hand, if $a_i = 1$ for $1 \leq i \leq n$ and $\mathcal{C} = \{c_1, \dots, c_\ell\}$ with $0 < c_1 \leq \dots \leq c_\ell$, then we have

$$\text{reg}(S) = \{m \in \mathbb{Z} : m \geq (n - 1)(c_\ell - 1)\}.$$

Example 4.3. Let $t \in \mathbb{N}$. Suppose that S is the homogeneous coordinate ring of the Hirzebruch surface \mathbb{F}_t ; see Example 2.6. If $\mathcal{C} = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ then Example 3.5 implies that

$$\text{reg}(S) = \begin{cases} \mathbb{N}^2 & \text{for } t = 0, 1, \\ \left(\begin{bmatrix} t-1 \\ 0 \end{bmatrix} + \mathbb{N}^2 \right) \cup \left(\begin{bmatrix} 0 \\ 1 \end{bmatrix} + \mathbb{N}^2 \right) & \text{for } t \geq 2. \end{cases}$$

Observe that when $t \geq 2$ the set $\text{reg}(S)$ is not determined by a single element of G . Moreover, we have $\text{int } \mathcal{K}^{\text{sat}} \subset \text{reg}(S)$ for all $t \in \mathbb{N}$, but $\mathbf{0} \notin \text{reg}(S)$ for $t \geq 2$. The shaded region in Figure 5 represents $\text{reg}(S)$ when $t = 2$.




Figure 5. The $\text{reg}(S)$ for the homogeneous coordinate ring S of \mathbb{F}_2 .

Without additional hypotheses, it is possible that $\text{reg}(M)$ is the empty set. Indeed, if $S = \mathbb{k}[x_1, x_2, x_3]$ has the \mathbb{Z} -grading described in Example 3.10, then $\text{reg}(S) = \emptyset$. Fortunately, there is a large class of pairs (S, B) for which $\text{reg}(M) \neq \emptyset$ for all finitely generated S -modules M . Specifically, we have the following:

Proposition 4.4. *If $\text{pos}(\mathcal{A})$ is a pointed cone and $\mathbb{N}\mathcal{C} \cap \text{int } \mathcal{K}^{\text{sat}} \neq \emptyset$, then every module M is \mathbf{m} -regular for some $\mathbf{m} \in G$.*

Proof. Corollary 3.8 states that there is $\mathbf{m} \in \mathcal{K}^{\text{sat}}$ such that $H_B^i(M)_{\mathbf{p}} = 0$ for all i and all $\mathbf{p} \in \mathbf{m} + \mathcal{K}^{\text{sat}}$. By hypothesis, there exists $\mathbf{c} \in \mathbb{N}\mathcal{C} \cap \text{int } \mathcal{K}^{\text{sat}}$. Remark 3.7 implies that there exists $k \in \mathbb{N}$ such that

$$k\mathbf{c} \in \bigcap_{\substack{\lambda_1, \dots, \lambda_\ell \in \mathbb{N} \\ \lambda_1 + \dots + \lambda_\ell \leq d}} (\mathbf{m} + \lambda_1 \mathbf{c}_1 + \dots + \lambda_\ell \mathbf{c}_\ell + \mathcal{K}^{\text{sat}}).$$

We conclude that $k\mathbf{c} \in \text{reg}(M)$. \square

To show that a module M is \mathbf{m} -regular directly from Definition 4.1, one must verify that an infinite number of graded components of the local cohomology vanishes. In fact, under the appropriate hypothesis, one need only check a finite number of components. To prove this result, we introduce the following two weaker versions of regularity.

Definition 4.5. Given $k \in \mathbb{N}$, the module M is \mathbf{m} -regular from level k if $H_B^i(M)_{\mathbf{p}} = 0$ for all $i \geq k$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$. In particular, M is \mathbf{m} -regular if and only if it is \mathbf{m} -regular from level 0. We set $\text{reg}^k(M) := \{\mathbf{m} \in G : M \text{ is } \mathbf{m}\text{-regular from level } k\}$. For

$k > 0$, we say M is weakly \mathbf{m} -regular from level k if $H_B^i(M)_{\mathbf{p}} = 0$ for all $i \geq k$ and all $\mathbf{p} = \mathbf{m} - \lambda_1 \mathbf{c}_1 - \cdots - \lambda_\ell \mathbf{c}_\ell$ where $\lambda_j \in \mathbb{N}$ and $\lambda_1 + \cdots + \lambda_\ell = i - 1$.

Our goal is to show that, when $\mathcal{C} \subseteq \mathcal{H}$, M is \mathbf{m} -regular if and only if it is weakly \mathbf{m} -regular from level 1 and $H_B^0(M)_{\mathbf{p}} = 0$ for $\mathbf{p} \in \bigcup_{1 \leq j \leq \ell} (\mathbf{m} + \mathbf{c}_j + \mathbb{N}\mathcal{C})$. To accomplish this, we need the following fact.

Lemma 4.6. *Let k be a positive integer. If M is weakly \mathbf{m} -regular from level k and $g \in S_{\mathbf{c}_j}$ is almost a nonzerodivisor on M for some $1 \leq j \leq \ell$, then $M/g_j M$ is also weakly \mathbf{m} -regular from level k .*

Proof. Since g_j is almost a nonzerodivisor on M , the submodule $(0 :_M g_j)$ is a B -torsion module. We set $\bar{M} := M/(0 :_M g_j)$. Because $H_B^i((0 :_M g_j)) = 0$ for $i > 0$, the long exact sequence associated to the exact sequence $0 \rightarrow (0 :_M g_j) \rightarrow M \rightarrow \bar{M} \rightarrow 0$ implies that $H_B^i(M) = H_B^i(\bar{M})$ for $i > 0$. Since M is weakly \mathbf{m} -regular from level k and $k > 0$, it follows that \bar{M} is weakly \mathbf{m} -regular from level k . On the other hand, the long exact sequence associated to the exact sequence $0 \rightarrow \bar{M}(-\mathbf{c}_j) \rightarrow M \rightarrow M/g_j M \rightarrow 0$ contains

$$(4.6.1) \quad H_B^i(M)_{\mathbf{p}} \rightarrow H_B^i(M/g_j M)_{\mathbf{p}} \rightarrow H_B^{i+1}(\bar{M})_{\mathbf{p}-\mathbf{c}_j}.$$

Since M and \bar{M} are weakly \mathbf{m} -regular from level k , the left and right modules in (4.6.1) vanish when $i \geq k$ and $\mathbf{p} = \mathbf{m} - \lambda_1 \mathbf{c}_1 - \cdots - \lambda_\ell \mathbf{c}_\ell$ where $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$ and

$$\lambda_1 + \cdots + \lambda_\ell = i - 1.$$

Therefore, the middle module also vanishes which proves that $M/g_j M$ is weakly \mathbf{m} -regular from level k . \square

Theorem 4.7. *Let k be a positive integer. If $\mathcal{C} \subseteq \mathcal{H}$ and M is weakly \mathbf{m} -regular from level k , then M is weakly \mathbf{p} -regular from level k for every $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$.*

Proof. Since extension of our base field commutes with the formation of local cohomology, we may assume for the proof that \mathbb{k} is infinite. Because $\mathcal{C} \subseteq \mathcal{H}$, Proposition 3.1 implies that we may choose a nonzerodivisor $g_j \in S_{\mathbf{c}_j}$ on M for each $1 \leq j \leq \ell$.

Suppose that $k > 0$ and that M is weakly \mathbf{m} -regular from level k . Since every $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$ can be expressed in the form $\mathbf{p} = \mathbf{m} + \lambda_1 \mathbf{c}_1 + \cdots + \lambda_\ell \mathbf{c}_\ell$ where $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$, it suffices to prove that if M is weakly \mathbf{q} -regular from level k then M is also weakly $(\mathbf{q} + \mathbf{c}_j)$ -regular from level k for each $1 \leq j \leq \ell$. We proceed by induction on $\dim M$. If $\dim M = 0$ then Grothendieck's vanishing theorem ([5], Theorem 6.1.2) implies that $H_B^i(M) = 0$ for $i > 0$. Thus, M is weakly \mathbf{q} -regular from level k for all $\mathbf{q} \in G$ and there is nothing more to prove.

Assume that $\dim M > 0$ and set $\bar{M} := M/H_B^0(M)$. Since $H_B^0(M)$ is a B -torsion module, the long exact sequence in cohomology arising from the short exact sequence $0 \rightarrow H_B^0(M) \rightarrow M \rightarrow \bar{M} \rightarrow 0$ implies that $H_B^i(M) = H_B^i(\bar{M})$ for all $i > 0$. Hence, \bar{M} is weakly \mathbf{q} -regular from level k and it suffices to show that \bar{M} is weakly $(\mathbf{q} + \mathbf{c}_j)$ -regular from level k .

Since g_j is a nonzerodivisor on \bar{M} , we have $\dim \bar{M}/g_j\bar{M} < \dim \bar{M}$. Lemma 4.6 shows that $\bar{M}/g_j\bar{M}$ is weakly \mathbf{q} -regular from level k and the induction hypothesis implies $\bar{M}/g_j\bar{M}$ is also weakly $(\mathbf{q} + \mathbf{c}_j)$ -regular from level k . Taking the long exact sequence associated to the exact sequence $0 \rightarrow \bar{M}(-\mathbf{c}_j) \rightarrow \bar{M} \rightarrow \bar{M}/g_j\bar{M} \rightarrow 0$, we obtain the exact sequence $H_B^i(\bar{M})_{\mathbf{q}'} \rightarrow H_B^i(\bar{M})_{\mathbf{q}'+\mathbf{c}_j} \rightarrow H_B^i(\bar{M}/g_j\bar{M})_{\mathbf{q}'+\mathbf{c}_j}$. Since \bar{M} is weakly \mathbf{q} -regular from level k and $\bar{M}/g_j\bar{M}$ is $(\mathbf{q} + \mathbf{c}_j)$ -regular from level k , the left and right modules vanish when $i \geq k$ and $\mathbf{q}' = \mathbf{q} - \lambda_1 \mathbf{c}_1 - \dots - \lambda_\ell \mathbf{c}_\ell$ where $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$ and $\lambda_1 + \dots + \lambda_\ell = i - 1$. Therefore, the middle module also vanishes which proves \bar{M} is weakly $(\mathbf{q} + \mathbf{c}_j)$ -regular from level k . \square

Theorem 4.7 provides the desired alternative characterization of \mathbf{m} -regularity.

Corollary 4.8. *Let $\mathcal{C} \subseteq \mathcal{K}$. The module M is \mathbf{m} -regular if and only if M is weakly \mathbf{m} -regular from level 1 and $H_B^0(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathbf{m} + \left(\bigcup_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathbb{N}\mathcal{C}) \right)$.*

Proof. Suppose that the module M is \mathbf{m} -regular from level 1 and $H_B^0(M)_{\mathbf{p}}$ vanishes for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1]$. Since the condition on $H_B^0(M)_{\mathbf{p}}$ is the same as in Definition 4.1, we only need to show that $H_B^i(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$. However, this is the content of Theorem 4.7. The converse follows from the Definition 4.1 and Definition 4.5 \square

The next example illustrates that the condition $\mathcal{C} \subseteq \mathcal{K}$ is necessary for Theorem 4.7.

Example 4.9. Let $S = \mathbb{k}[x_1, x_2]$ have the \mathbb{Z} -grading defined by $\deg(x_1) = 2$ and $\deg(x_2) = 3$ and let $B = \langle x_1, x_2 \rangle$. Example 2.1 shows that $\mathcal{K} = 6\mathbb{N}$. Proposition 3.2 establishes that the nonvanishing local cohomology is concentrated in $H_B^2(S)$ and that $H_B^2(S)$ is nonzero exactly in degrees $-5, -7, -8, -9, \dots$. If $\mathcal{C} = \{\mathbf{1}\} \not\subseteq \mathcal{K}$, then S is weakly (-5) -regular from level 1 but not weakly (-4) -regular from level 1. The strategy used in the proof of Theorem 4.7 does not apply because $S_1 = \emptyset$, so there is no nonzerodivisor of degree 1.

By design, the regularity of a module measures where its cohomological complexities vanish. Since Gröbner bases calculations are linked to homological properties, one expects a strong connection between regularity and computational complexity. Both Proposition 6.13 and Corollary 7.3 provide further support for this idea by relating regularity to initial modules and free resolutions. To make the connection between regularity and complexity more precise, we are interested in the following open problem:

Problem 4.10. Give a computationally efficient method of calculating $\text{reg}(M)$.

When S has the standard grading, this problem is solved in [1] by showing that $\text{reg}(M)$ is determined by the largest degree generator of the initial module $\text{in}(M)$ with respect to a reverse lexicographic order in generic coordinates. Unfortunately, this technique does not extend directly to polynomial rings with arbitrary multigradings. As the next example demonstrates, there may not be any coordinate changes which preserve the grading and change the initial module.

Example 4.11. Let S be the homogeneous coordinate ring of a toric variety X obtained from \mathbb{P}^2 by a sequence of five blow-ups. More explicitly, the minimal lat-

tice points \mathcal{B} on the rays of the fan correspond to the columns of the matrix $\begin{bmatrix} 1 & 1 & 0 & -1 & -1 & -1 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 & -1 & -1 & -1 \end{bmatrix}$ and the associated irrelevant ideal is

$$B = \langle x_3x_4x_5x_6x_7x_8, x_1x_4x_5x_6x_7x_8, x_1x_2x_5x_6x_7x_8, x_1x_2x_3x_6x_7x_8, \\ x_1x_2x_3x_4x_7x_8, x_1x_2x_3x_4x_5x_8, x_2x_3x_4x_5x_6x_7, x_1x_2x_3x_4x_5x_6 \rangle.$$

Hence, $G = \mathbb{Z}^6$ and we may assume that \mathcal{A} be given by the columns of the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & -2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -2 & 1 \end{bmatrix}.$$

The homogeneous coordinate ring $S = \mathbb{k}[x_1, \dots, x_8]$ has the \mathbb{Z}^6 -grading induced by $\deg(x_i) = \mathbf{a}_i$ for $1 \leq i \leq 8$. Since $S_{\mathbf{a}_i}$ is the \mathbb{k} -span of the variable x_i for $1 \leq i \leq 8$, Corollary 4.7 in [11] establishes that $\text{Aut}(S) = (\mathbb{k}^*)^8$. As a consequence, any change of coordinates which preserves the grading does not alter the initial ideal. Hence, one cannot develop a theory of generic initial ideals.

5. Degrees of generators

The regularity of a module should be regarded as a measure of its complexity. In this section, we justify this idea by proving that the regularity controls the degrees of the minimal generators. To understand the minimal generators of a module, we study submodules of the following form.

Definition 5.1. Let \mathcal{D} be a subset of G . We define the \mathcal{D} -truncation of M , denoted $M|_{\mathcal{D}}$, to be the submodule of M generated by all the homogeneous elements in M of degree \mathbf{p} where $\mathbf{p} \in \mathcal{D}$.

In contrast with the standard graded case, the next example illustrates that $(M|_{\mathcal{D}})_{\mathbf{p}}$ may be nonzero even if $\mathbf{p} \notin \mathcal{D} + \mathbb{N}\mathcal{H}^{\text{sat}}$.

Example 5.2. Suppose that S is the homogeneous coordinate ring of the Hirzebruch surface \mathbb{F}_2 ; see Examples 2.2 and 2.6. If $\mathcal{D} := \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \mathbb{N}^2$, then $S|_{\mathcal{D}}$ is generated in degree $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Observe that $x_1^3x_2^2 \in S|_{\mathcal{D}}$ but $\deg(x_1^3x_2^2) \notin \mathcal{D}$.

To prove the main theorem in this section, we need the following fact.

Lemma 5.3. Let $\mathbf{c} \in \mathbb{N}\mathcal{C}$ and let $\mathcal{V} \subseteq G$ such that $H_B^0(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathcal{V}$. If M is \mathbf{m} -regular from level 1 and $g \in S_{\mathbf{c}}$ is almost a nonzerodivisor on M , then $H_B^0(M/gM)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathcal{V} \cap (\mathbf{m} + \mathbf{c} + \mathbb{N}\mathcal{C})$.

Proof. Set $\bar{M} := M/(0 :_M g)$. Because g is almost a nonzerodivisor on M , $(0 :_M g)$ is a B -torsion module and the long exact sequence associated to the short exact sequence $0 \rightarrow (0 :_M g) \rightarrow M \rightarrow \bar{M} \rightarrow 0$ implies that $H_B^i(M) = H_B^i(\bar{M})$ for $i > 0$. Since M is \mathbf{m} -regular from level 1, it follows that \bar{M} is also \mathbf{m} -regular from level 1. Now, the long exact sequence associated to the exact sequence $0 \rightarrow \bar{M}(-\mathbf{c}) \rightarrow M \rightarrow M/gM \rightarrow 0$ contains $H_B^0(M)_\mathbf{p} \rightarrow H_B^0(M/gM)_\mathbf{p} \rightarrow H_B^1(\bar{M})_{\mathbf{p}-\mathbf{c}}$. By hypothesis, the left module vanishes when $\mathbf{p} \in \mathcal{V}$. Since \bar{M} is \mathbf{m} -regular from level 1, the right module vanishes for all $\mathbf{p} \in \mathbf{m} + \mathbf{c} + \mathbb{N}\mathcal{C}$. Hence, the middle module vanishes for all $\mathbf{p} \in \mathcal{V} \cap (\mathbf{m} + \mathbf{c} + \mathbb{N}\mathcal{C})$. \square

We now prove that if M is \mathbf{m} -regular then $M|_{\mathbf{m}} = M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$. In the standard graded case, this is true for any $\mathbf{m} \in \mathbb{Z}$ that is larger than the maximum degree of the minimal generators.

Theorem 5.4. *Assume that $\mathcal{C} \subseteq \mathcal{K}$. If the module M is \mathbf{m} -regular, then we have $M|_{\mathbf{m}} = M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$. In particular, the degrees of the minimal generators of M do not belong to the set $\text{reg}(M) + \left(\bigcup_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathbb{N}\mathcal{C}) \right)$.*

Proof. We prove the following claim: If M is \mathbf{m} -regular from level 1 and $H_B^0(M)_\mathbf{p} = 0$ for all $\mathbf{p} \in \mathcal{V}$, then $M|_{(\mathbf{m} + \mathbb{N}\mathcal{C}) \cap \mathcal{V}}$ is a submodule of $M|_{\mathbf{m}}$. We proceed by induction on $\dim M$. If $\dim M < 0$, which is equivalent to saying that $M = 0$, then $M|_{(\mathbf{m} + \mathbb{N}\mathcal{C}) \cap \mathcal{V}} = 0$ is trivially a submodule of $M|_{\mathbf{m}}$.

Suppose that $\dim M \geq 0$. Set $\bar{M} := M/H_B^0(M)$ and consider the short exact sequence

$$(5.4.1) \quad 0 \rightarrow H_B^0(M) \rightarrow M \rightarrow \bar{M} \rightarrow 0.$$

We claim that it is enough to prove that $\bar{M}|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$ is a submodule of $\bar{M}|_{\mathbf{m}}$. To see this, let $\mathbf{p} \in (\mathbf{m} + \mathbb{N}\mathcal{C}) \cap \mathcal{V}$ and let $f \in M_\mathbf{p}$. If $\bar{f} \in \bar{M}_\mathbf{p}$ can be written as $\sum_i s_i \bar{f}_i$ for some $s_i \in S$ and some $\bar{f}_i \in \bar{M}_\mathbf{m}$, then $f = \sum_i s_i f_i + h$ for $h \in H_B^0(M)_\mathbf{p}$. Since $\mathbf{p} \in \mathcal{V}$, we have $H_B^0(M)_\mathbf{p} = 0$ which implies $h = 0$. Thus, $f = \sum_i s_i f_i$ and it suffices to prove the claim for \bar{M} . The long exact sequence associated to (5.4.1) implies $H_B^i(M) = H_B^i(\bar{M})$ for all $i > 0$. Since M is \mathbf{m} -regular from level 1 and $H_B^0(\bar{M}) = 0$, we deduce that \bar{M} is \mathbf{m} -regular.

Extending the base field commutes with the computing local cohomology, so we may assume without loss of generality that \mathbb{k} is infinite. Because $\mathcal{C} \subseteq \mathcal{K}$, Proposition 3.1 implies that, for each $1 \leq j \leq \ell$, we may choose a nonzerodivisor $g_j \in S_{\mathbf{c}_j}$ on \bar{M} . It follows from Lemma 4.6 that $\bar{M}/g_j \bar{M}$ is \mathbf{m} -regular from level 1 and Lemma 5.3 shows that $H_B^0(\bar{M}/g_j \bar{M})_\mathbf{p} = 0$ for all $\mathbf{p} \in \mathbf{m} + \mathbf{c}_j + \mathbb{N}\mathcal{C}$. Since $\dim \bar{M}/g_j \bar{M} < \dim \bar{M} \leq \dim M$, the induction hypothesis with $\mathcal{V} = \mathbf{m} + \mathbf{c}_j + \mathbb{N}\mathcal{C}$ guarantees that $(\bar{M}/g_j \bar{M})|_{(\mathbf{m} + \mathbf{c}_j + \mathbb{N}\mathcal{C})}$ is a submodule of $(\bar{M}/g_j \bar{M})|_{\mathbf{m}}$. It remains to show that this implies $\bar{M}|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$ is a submodule of $\bar{M}|_{\mathbf{m}}$.

Suppose otherwise. For each element $f \in \bar{M}$ with $\deg(f) \in \mathbf{m} + \mathbb{N}\mathcal{C}$, we set

$$\text{reach}(f) := \min\{\lambda_1 + \cdots + \lambda_\ell : \deg(f) = \mathbf{m} + \lambda_1 \mathbf{c}_1 + \cdots + \lambda_\ell \mathbf{c}_\ell \text{ and } \lambda_1, \dots, \lambda_\ell \in \mathbb{N}\}.$$

Because \bar{M} is noetherian and we are assuming that $\bar{M}|_{(m+\mathbb{N}\mathcal{C})} \not\subseteq \bar{M}|_m$, there is a minimal generator $f \in \bar{M}|_{(m+\mathbb{N}\mathcal{C})}$ which has smallest $\text{reach}(f)$ among all the minimal generators of $\bar{M}|_{(m+\mathbb{N}\mathcal{C})}$ with degree not equal to m . Since $f \notin \bar{M}|_m$, we have $\text{reach}(f) > 0$ and there exists c_j with $1 \leq j \leq \ell$ such that $\text{deg}(f) \in m + c_j + \mathbb{N}\mathcal{C}$. From the previous paragraph, we know that the image of f in $(\bar{M}/g_j\bar{M})|_{(m+c_j+\mathbb{N}\mathcal{C})}$ belongs to $(\bar{M}/g_j\bar{M})|_m$. Hence, we may choose homogeneous elements $f_1, \dots, f_e \in \bar{M}|_m$ and $s_1, \dots, s_e \in S$ such that

$$f - s_1f_1 - \dots - s_ef_e \in (g_j\bar{M})|_{(m+c_j+\mathbb{N}\mathcal{C})} \subseteq g_j(\bar{M}|_{(m+\mathbb{N}\mathcal{C})}).$$

Let f' be the homogeneous element of $\bar{M}|_{(m+\mathbb{N}\mathcal{C})}$ satisfying the equation

$$f - s_1f_1 - \dots - s_ef_e = g_jf'.$$

Since f does not belong to $\bar{M}|_m$, the element f' cannot belong to $\bar{M}|_m$. However, $\text{reach}(f') < \text{reach}(f)$ and $f' \in \bar{M}|_{(m+\mathbb{N}\mathcal{C})}$ which contradicts our choice of f . We conclude that $\bar{M}|_{(m+\mathbb{N}\mathcal{C})}$ is a submodule of $\bar{M}|_m$. \square

Proof of Theorem 1.3. In the introduction we assumed that \mathcal{C} was the minimal generating set of \mathcal{K} . Hence, Theorem 1.3 follows from Theorem 5.4. \square

When S has the standard grading and $\mathcal{C} = \{1\}$, the statement that the minimal generators of M do not lie in $m + \mathbb{N}\mathcal{C}[1]$ is equivalent to saying that the minimal generators of M have degree at most m . In this case, Theorem 5.4 proves that the regularity gives a bound on the degrees of the minimal generators. For an ideal in S , an upper bound on the minimal generators yields a finite set containing the degrees of the minimal generators. In contrast, Theorem 5.4 does not automatically produce a finite set containing the degrees of the minimal generators for an ideal when S has a general multigrading.

We may still ask the question whether if M is m -regular all minimal generators of M lie in $m - \mathbb{N}\mathcal{C}$. It is not hard to find examples showing that this need not be the case, but to date all such examples have had $H_B^0(M) \neq 0$. In the bigraded case Hoffman and Wang [25] take a different approach, adding an extra local cohomology vanishing requirement to guarantee this bound on generators.

6. Regularity of \mathcal{O}_X -modules

In this section, we develop a multigraded version of regularity for coherent sheaves on a simplicial toric variety X . Since \mathcal{O}_X -modules correspond to finitely generated modules over the homogeneous coordinate ring of X , regularity for \mathcal{O}_X -modules is essentially a special case of regularity for S -modules. Nevertheless, the geometric context provides new interpretations and applications for regularity.

To begin, we examine the relation between sheaves on a toric variety X and G -graded modules over its homogeneous coordinate ring S ; see [11]. For $\sigma \in \Delta$, we write x_σ for the monomial $\prod_{i \notin \sigma} x_i \in S$ and $U_\sigma \cong \text{Spec}(S[x_\sigma^{-1}]_0)$ is the corresponding open affine subset of X . Every (not necessarily finitely generated) G -graded S -module F gives rise to a quasicoherent sheaf \tilde{F} on X where $H^0(U_\sigma, \tilde{F}) = (F[x_\sigma^{-1}])_0$. If M is finitely generated G -graded S -

module, then \tilde{M} is a coherent \mathcal{O}_X -module. Moreover, every quasicoherent sheaf on X is of this form for some G -graded S -module F and if the sheaf is coherent then F can be taken to be finitely generated. For $\mathbf{p} \in G$, the sheaf associated to the module $S(\mathbf{p})$ is denoted $\mathcal{O}_X(\mathbf{p})$. Note that the natural map $\mathcal{O}_X(\mathbf{p}) \otimes_{\mathcal{O}_X} \mathcal{O}_X(\mathbf{q}) \rightarrow \mathcal{O}_X(\mathbf{p} + \mathbf{q})$ need not be an isomorphism. For an \mathcal{O}_X -module \mathcal{F} , we simply write $\mathcal{F}(\mathbf{p})$ for $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_X(\mathbf{p})$.

The map sending a finitely generated G -graded S -module M to the sheaf \tilde{M} is not injective. In fact, there are many nonzero modules which give the zero sheaf. The following proposition analyzes this phenomenon.

Proposition 6.1. *The sheaf \tilde{M} is zero if and only if there is $j > 0$ such that $B^j M_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathcal{K}$.*

Proof. Suppose that $B^j M_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathcal{K}$. Fix $\sigma \in \Delta$ and take $f/x^u \in (M[x_{\hat{\sigma}}^{-1}])_0$. Remark 3.7 implies that there exists a monomial $\mathbf{x}^v \in S$ such that $\text{supp}(v) \subseteq \hat{\sigma}$ and $\deg(\mathbf{x}^{u+v}) \in \mathcal{K}^{\text{sat}}$. Moreover, we may choose $k \in \mathbb{N}$ such that $\mathbf{p} := \deg(\mathbf{x}^{ku+kv}) \in \mathcal{K}$. It follows that $\mathbf{x}^{(k-1)u+kv} f \in M_{\mathbf{p}}$ and $(x_{\hat{\sigma}})^j \mathbf{x}^{(k-1)u+kv} f = 0$ for some $j > 0$. Therefore, $f/x^u = 0$ and $(M[x_{\hat{\sigma}}^{-1}])_0 = 0$ which shows that \tilde{M} is the zero sheaf.

Conversely, suppose that \tilde{M} is the zero sheaf, fix $\mathbf{p} \in \mathcal{K}$ and let $f \in M_{\mathbf{p}}$. Lemma 2.4 implies that for each $\sigma \in \Delta$ there is a monomial $\mathbf{x}^u \in S$ such that $\deg(\mathbf{x}^u) = \mathbf{p}$ and $\text{supp}(u) \subseteq \hat{\sigma}$. The monomial \mathbf{x}^u is invertible in $S[x_{\hat{\sigma}}^{-1}]$ and $f/x^u \in (M[x_{\hat{\sigma}}^{-1}])_0 = 0$. It follows that $(x_{\hat{\sigma}})^j f = 0$ for some $j > 0$. To see that one j works for all $f \in M_{\mathbf{p}}$ and all $\mathbf{p} \in \mathcal{K}$, observe that $M|_{\mathcal{X}}$ is a submodule of M and thus finitely generated. The proposition now follows easily. \square

As above, we continue to write $\mathcal{C} = \{c_1, \dots, c_\ell\}$ for a fixed subset of G , the class group of X . Geometrically, the set \mathcal{C} corresponds to choosing a collection of reflexive sheaves on X . When X is smooth, $G = \text{Pic}(X)$ and \mathcal{C} corresponds to a choice of line bundles. In particular, fixing an ample line bundle on X determines a set \mathcal{C} consisting of one element. At the other extreme, the toric variety X is equipped with a canonical choice for \mathcal{C} , namely the minimal generators of the semigroup \mathcal{K} . We are most interested in this case. Nonetheless, we expect that the flexibility in allowing other sets \mathcal{C} will also be useful in studying how maps between toric varieties effect regularity.

In analogy with Definition 4.1, we have the following:

Definition 6.2. If $\mathbf{m} \in G$, then an \mathcal{O}_X -module \mathcal{F} is *\mathbf{m} -regular* (with respect to \mathcal{C}) if $H^i(X, \mathcal{F}(\mathbf{p})) = 0$ for all $i > 0$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[-i]$. We write $\text{reg}(\mathcal{F})$ for the set $\{\mathbf{m} \in G : \mathcal{F} \text{ is } \mathbf{m}\text{-regular}\}$.

Remark 6.3. Notice that if \mathcal{F} is \mathbf{m} -regular then \mathcal{F} is \mathbf{p} -regular for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$.

To make an explicit connection between the two definitions of regularity, we first recall the following. If \mathcal{F} is the coherent \mathcal{O}_X -module corresponding to M , then the local cohomology is related to the (Zariski) cohomology of sheaves by the exact sequence

$$(6.3.1) \quad 0 \rightarrow H_B^0(M) \rightarrow M \rightarrow \bigoplus_{\mathbf{p} \in G} H^0(X, \mathcal{F}(\mathbf{p})) \rightarrow H_B^1(M) \rightarrow 0$$

and $H_B^{i+1}(M) \cong \bigoplus_{\mathbf{p} \in G} H^i(X, \mathcal{F}(\mathbf{p}))$ for all $i \geq 1$; see [16], Proposition 2.3. Using this observation, we have:

Proposition 6.4. *Let \mathcal{F} be the coherent \mathcal{O}_X -module associated to the module M . The sheaf \mathcal{F} is \mathbf{m} -regular if and only if M is \mathbf{m} -regular from level 2. The module M is \mathbf{m} -regular if and only if the following three conditions hold:*

- (1) \mathcal{F} is \mathbf{m} -regular.
- (2) The natural map $M_{\mathbf{p}} \rightarrow H^0(X, \mathcal{F}(\mathbf{p}))$ is surjective for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$.
- (3) $H_B^0(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \bigcup_{1 \leq j \leq \ell} (\mathbf{m} + \mathbf{c}_j + \mathbb{N}\mathcal{C})$.

Proof. Since $H_B^{i+1}(M)_{\mathbf{p}} \cong H^i(X, \mathcal{F}(\mathbf{p}))$ for all $i \geq 1$, we see that \mathcal{F} is \mathbf{m} -regular if and only if M is \mathbf{m} -regular from level 2. To establish the second part, observe that saying M is \mathbf{m} -regular is equivalent to saying that M is \mathbf{m} -regular from level 2, $H_B^1(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$ and $H_B^0(M)_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathbf{m} + \bigcup_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathbb{N}\mathcal{C})$. Since the exact sequence (6.3.1) implies that $H_B^1(M)_{\mathbf{p}} = 0$ if and only if $M_{\mathbf{p}} \rightarrow H^0(X, \mathcal{F}(\mathbf{p}))$ is surjective, the assertion follows. \square

Example 6.5. Since $H_B^i(S) = 0$ for $i = 0, 1$ (see Remark 3.3), Proposition 6.4 implies that $\text{reg}(\mathcal{O}_X) = \text{reg}(S)$. In particular, Example 4.2 shows that $\text{reg}(\mathcal{O}_{\mathbb{P}^d}) = \mathbb{N}$ when $\mathcal{C} = \{\mathbf{1}\}$.

The next corollary shows that Definition 6.2 extends the original definition given in [30], §14. When $X = \mathbb{P}^d$ and $\mathcal{C} = \{\mathbf{1}\}$ ($\mathbf{1} \in G$ corresponds to $\mathcal{O}_X(\mathbf{1}) \in \text{Pic}(\mathbb{P}^d)$), the second part of this corollary is Mumford's definition.

Corollary 6.6. *Let \mathcal{F} be the coherent \mathcal{O}_X -module associated to the module M . If $\mathcal{C} \subseteq \mathcal{K}$ then the following are equivalent:*

- (1) \mathcal{F} is \mathbf{m} -regular.
- (2) $H^i(X, \mathcal{F}(\mathbf{m} - \lambda_1 \mathbf{c}_1 - \cdots - \lambda_\ell \mathbf{c}_\ell)) = 0$ for all $i > 0$ and all $\lambda_1, \dots, \lambda_\ell \in \mathbb{N}$ satisfying $\lambda_1 + \cdots + \lambda_\ell = i$.

Proof. This follows from Proposition 6.4 and Corollary 4.8. \square

The regularity of a finite set of points has a geometric interpretation. [28], Example 2.16 rephrases the following result in terms of the associated multigraded Hilbert polynomial.

Proposition 6.7. *Assume that $\mathcal{C} \subseteq \mathcal{K}$. Let Y be an artinian subscheme of X of length t (for example, a set of t points in X) and let I_Y be the associated B -saturated ideal in S . If $R_Y = S/I_Y$ then $\mathbf{m} \in \text{reg}(R_Y)$ if and only if the space of forms vanishing on Y has codimension t is the space of forms of degree \mathbf{m} .*

Proof. Since I_Y is B -saturated, the local cohomology module $H_B^0(R_Y)$ vanishes.

Because $\dim Y = 0$, we also have $H_B^{i+1}(R_Y) = \bigoplus_{\mathbf{p} \in G} H^i(X, \mathcal{O}_Y(\mathbf{p})) = 0$ for all $i > 0$. Hence, $\text{reg}(R_Y)$ is determined by the module $H_B^1(R_Y)$. Since the exact sequence (6.3.1) becomes $0 \rightarrow R_Y \rightarrow \bigoplus_{\mathbf{p} \in G} H^0(X, \mathcal{O}_Y(\mathbf{p})) \rightarrow H_B^1(R_Y) \rightarrow 0$, we see that $\mathbf{m} \in \text{reg}(R_Y)$ if and only if $(R_Y)_{\mathbf{m}} \rightarrow \bigoplus_{\mathbf{p} \in G} H^0(X, \mathcal{O}_Y(\mathbf{p}))$ is surjective. Again, because $\dim Y = 0$, the scheme Y is an affine variety and every reflexive sheaf on Y is trivial. It follows that for all \mathbf{p} we have $\mathcal{O}_Y(\mathbf{p}) \cong \mathcal{O}_Y$ and $\dim_{\mathbb{k}} H^0(X, \mathcal{O}_Y) = t$. Therefore, R_Y is \mathbf{m} -regular if and only if $\dim_{\mathbb{k}}(R_Y)_{\mathbf{m}} = t$. In other words, the space of forms $(I_Y)_{\mathbf{m}}$ that vanish on Y has codimension t . \square

Although the definition of regularity may seem rather technical (“apparently silly” in Mumford’s words), the following theorem provides a geometric interpretation for regularity. Concretely, the regularity of a coherent sheaf measures how much one has to twist for the sheaf to be generated by its global sections. We first show that the module M and certain truncations of M give rise to the same sheaf.

Lemma 6.8. *Assume that $\mathcal{C} \subseteq \mathcal{K}$ and $\dim_{\mathbb{R}} \text{pos}(\overline{\mathcal{C}}) = r$. If $\mathbf{m} \in \mathbb{Z}\mathcal{K}$ and M' is the quotient module $M/M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$ then there exists $j > 0$ such that $B^j M'_{\mathbf{p}} = 0$ for all $\mathbf{p} \in \mathbb{Z}\mathcal{K}$. Moreover, the module M and its truncation $M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$ correspond to the same \mathcal{O}_X -module.*

Proof. Fix $\mathbf{p} \in \mathbb{Z}\mathcal{K}$ and $f \in M_{\mathbf{p}}$. If $\mathbf{c} \in \text{int}\mathcal{K}$, then Remark 3.7 (applied to the group $\mathbb{Z}\mathcal{K}$) shows that there is $k \in \mathbb{N}$ such that $\mathbf{p} + k\mathbf{c} \in \mathbf{m} + \mathcal{K}$. Since $\mathcal{C} \subseteq \mathcal{K}$ and $\dim_{\mathbb{R}} \text{pos}(\overline{\mathcal{C}}) = r$, there exists $\mathbf{q} \in \mathcal{K}$ for which $\mathbf{p} + k\mathbf{c} + \mathbf{q} \in \mathbf{m} + \mathbb{N}\mathcal{C}$. Now, Lemma 2.4 implies that $B \subseteq \sqrt{\langle S_{k\mathbf{c} + \mathbf{q}} \rangle}$ which means that there is $j > 0$ such that $B^j \subseteq \langle S_{k\mathbf{c} + \mathbf{q}} \rangle$. Hence, we have $B^j f \in M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}$. Because M is finitely generated, there is one j that works for all $f \in M_{\mathbf{p}}$ and all $\mathbf{p} \in \mathcal{K}$ which establishes the first part.

For the second part, consider the short exact sequence

$$(6.8.2) \quad 0 \rightarrow M|_{(\mathbf{m} + \mathbb{N}\mathcal{C})} \rightarrow M \rightarrow M' \rightarrow 0.$$

Since the functor $F \mapsto \tilde{F}$ is exact, it suffices to show that $\tilde{M}' = 0$. By Proposition 6.1, this is equivalent to the first part of the lemma. \square

Theorem 6.9. *Assume that $\mathcal{C} \subseteq \mathcal{K}$ and $\dim_{\mathbb{R}} \text{pos}(\overline{\mathcal{C}}) = r$. If $\mathbf{m} \in \mathbb{Z}\mathcal{K}$ and the sheaf \mathcal{F} is \mathbf{m} -regular then the natural map*

$$(6.9.3) \quad H^0(X, \mathcal{F}(\mathbf{p})) \otimes H^0(X, \mathcal{O}_X(\mathbf{q})) \rightarrow H^0(X, \mathcal{F}(\mathbf{p} + \mathbf{q}))$$

is surjective for all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}$ and all $\mathbf{q} \in \mathbb{N}\mathcal{C}$. In particular, the sheaf $\mathcal{F}(\mathbf{p})$ is generated by its global sections.

Proof. Consider the G -graded S -module

$$M := \left(\bigoplus_{\mathbf{p} \in G} H^0(X, \mathcal{F}(\mathbf{p})) \right) \Big|_{(\mathbf{m} + \mathbb{N}\mathcal{C})}.$$

Because \mathcal{F} is the sheaf associated with the S -module $\bigoplus_{\mathbf{p} \in \mathcal{K}} H^0(X, \mathcal{F}(\mathbf{p}))$, Lemma 6.8 guarantees that $\tilde{M} = \mathcal{F}$. Since \mathcal{F} is \mathbf{m} -regular, Proposition 6.4 shows that M is \mathbf{m} -regular

from level 2. Combining the definition of M with the exact sequence (6.3.1), we have $H_B^i(M)_p = 0$ for all $i = 0, 1$ and all $p \in \mathfrak{m} + \mathbb{N}\mathcal{C}$. Hence, M is \mathfrak{m} -regular which implies that M is also p -regular for all $p \in \mathfrak{m} + \mathbb{N}\mathcal{C}$. If $p \in \mathfrak{m} + \mathbb{N}\mathcal{C}$, then Theorem 5.4 shows that $M|_p = M|_{(p+\mathbb{N}\mathcal{C})}$. In other words, $M_p \cdot S_q = M_{p+q}$ for all $q \in \mathbb{N}\mathcal{C}$ and this is equivalent to (6.9.3) being surjective. Furthermore, Lemma 6.8 also guarantees that \mathcal{F} is the sheaf associated to $M|_{(p+\mathbb{N}\mathcal{C})}$. Since $M|_p = M|_{(p+\mathbb{N}\mathcal{C})}$, this establishes the last part of the theorem. \square

Proof of Theorem 1.4. Part (1) follows from Remark 6.3. In the introduction, \mathcal{C} is the minimal generating set for \mathcal{K} . Therefore, Theorem 6.9 proves Parts (2) and (3). \square

When $\mathcal{F} = \mathcal{O}_X$, Theorem 6.9 has a classical geometric interpretation. Specifically, if $p \in \text{reg}(\mathcal{O}_X)$ and $\mathcal{O}_X(p)$ is an ample line bundle then Theorem 6.9 implies that the natural map $H^0(X, \mathcal{O}_X(p)) \otimes H^0(X, \mathcal{O}_X(jp)) \rightarrow H^0(X, \mathcal{O}_X((j+1)p))$ is surjective for all $j \geq 0$. It follows (see [23], Exercise II.5.14) that $\mathcal{O}_X(p)$ embeds X into \mathbb{P}^N as a projectively normal variety where $N := \dim_{\mathbb{k}} H^0(X, \mathcal{O}_X(p))$.

It is an interesting open problem to describe the set of ample line bundles on a d -dimensional toric variety X which give a projectively normal embedding. When $\mathcal{O}_X(\mathfrak{m})$ is an ample line bundle, Bruns, Gubeladze, and Trung [6] show that the complete linear series of the line bundle $\mathcal{O}_X((d-1)\mathfrak{m})$ produces a projectively normal embedding of X . Moreover, this bound is sharp on some simplicial toric varieties. For smooth toric varieties, the question of whether the map

$$H^0(X, \mathcal{O}_X(p)) \otimes H^0(X, \mathcal{O}_X(q)) \rightarrow H^0(X, \mathcal{O}_X(p+q))$$

is surjective for $\mathcal{O}_X(p)$ an ample line bundle and $\mathcal{O}_X(q)$ a nef line bundle appears in [32]. This would imply that every ample line bundle induced a projectively normal embedding.

These questions motivate the study of when $\mathbf{0} \in \text{reg}(\mathcal{O}_X)$.

Proposition 6.10. *If X is a finite product of projective spaces and \mathcal{C} is the set of unique minimal generators of \mathcal{K} , then the structure sheaf \mathcal{O}_X is $\mathbf{0}$ -regular.*

Proof. We proceed by induction on the number of factors in X . When there is only one copy of projective space, Example 6.5 establishes the proposition. For the induction step, we prove the following stronger claim: Let X be a smooth projective toric variety such that \mathcal{O}_X is $\mathbf{0}$ -regular with respect to $\{c_1, \dots, c_\ell\}$. If $Z = X \times \mathbb{P}^h$ then \mathcal{O}_Z is $\mathbf{0}$ -regular with respect to $\left\{ \begin{bmatrix} c_1 \\ \mathbf{0} \end{bmatrix}, \dots, \begin{bmatrix} c_\ell \\ \mathbf{0} \end{bmatrix}, \begin{bmatrix} \mathbf{0} \\ \mathbf{1} \end{bmatrix} \right\}$.

To accomplish this, we first fix some notation. Let Δ and Σ be the simplicial complexes arising from the fans of X and Z respectively. If the set $\{a_1, \dots, a_n\}$ defines the \mathbb{Z}^r -grading on the homogeneous coordinate ring of X then the homogeneous coordinate ring $S = \mathbb{k}[x_1, \dots, x_n, y_0, \dots, y_h]$ has the \mathbb{Z}^{r+1} -grading induced by $\deg(x_i) = \begin{bmatrix} a_i \\ \mathbf{0} \end{bmatrix}$ and $\deg(y_i) = \begin{bmatrix} \mathbf{0} \\ \mathbf{1} \end{bmatrix}$. For $\tau \subseteq [n]$ and $\sigma \subseteq \{0, \dots, h\}$, we write $\mathcal{V}_{\tau \cup \sigma}$ for the subset of \mathbb{Z}^{r+1} given by

$$\begin{aligned} & \left[\begin{array}{c} \sum_{j \in \tau} -\mathbf{a}_j \\ -|\sigma| \end{array} \right] + \mathbb{N} \left\{ \left[\begin{array}{c} \mathbf{a}_j \\ \mathbf{0} \end{array} \right] : j \in \hat{\tau} \right\} + \mathbb{N} \left\{ \left[\begin{array}{c} -\mathbf{a}_j \\ \mathbf{0} \end{array} \right] : j \in \tau \right\} \\ & + \mathbb{N} \left\{ \left[\begin{array}{c} \mathbf{0} \\ \mathbf{1} \end{array} \right] : \text{if } |\sigma| \neq h + 1 \right\} + \mathbb{N} \left\{ \left[\begin{array}{c} \mathbf{0} \\ -\mathbf{1} \end{array} \right] : \text{if } |\sigma| \neq 0 \right\}. \end{aligned}$$

We also set

$$\mathcal{R}^i := \left\{ -\lambda_0 \begin{bmatrix} \mathbf{0} \\ \mathbf{1} \end{bmatrix} - \lambda_1 \begin{bmatrix} \mathbf{c}_1 \\ \mathbf{0} \end{bmatrix} - \cdots - \lambda_\ell \begin{bmatrix} \mathbf{c}_\ell \\ \mathbf{0} \end{bmatrix} : \lambda_0, \dots, \lambda_\ell \in \mathbb{N} \text{ and } \lambda_0 + \cdots + \lambda_\ell = i \right\}.$$

Now, Corollary 3.4 implies that \mathcal{O}_Z is $\mathbf{0}$ -regular if $\mathcal{V}_{\tau \cup \sigma} \cap \mathcal{R}^i = \emptyset$ whenever $i > 0$ and $\tilde{H}^{i-1}(\Sigma_{\tau \cup \sigma}) \neq 0$. We consider four possible cases:

- If $\emptyset \neq \sigma \neq \{0, \dots, h\}$, then every maximal simplex in $\Sigma_{\tau \cup \sigma}$ contains σ . Since the induced subcomplex associated to σ is the standard simplex, it follows that $\Sigma_{\tau \cup \sigma}$ is contractible and $\tilde{H}^{i-1}(\Sigma_{\tau \cup \sigma}) = 0$.
- If $\sigma = \emptyset$ then $\Sigma_{\tau \cup \sigma}$ is a subcomplex of Δ . Since \mathcal{O}_X is $\mathbf{0}$ -regular, we must have $\mathcal{V}_{\tau \cup \sigma} \cap \mathcal{R}^i = \emptyset$ whenever $i > 0$ and $\tilde{H}^{i-1}(\Sigma_{\tau \cup \sigma}) \neq 0$.
- If $\sigma = \{0, \dots, h\}$ and $\tau \neq [n]$, then Alexander duality implies that $\tilde{H}^{i-1}(\Sigma_{\tau \cup \sigma})$ is isomorphic to $\tilde{H}^{d+h-i-1}(\Sigma_{\widehat{\tau \cup \sigma}})$ which reduces us to the second case.
- If $\sigma = \{0, \dots, h\}$ and $\tau = [n]$ then $\tilde{H}^{i-1}(\Sigma_{\tau \cup \sigma}) \neq 0$ if and only if $i = d + h$. Since both $\mathcal{O}_{\mathbb{P}^d}$ and \mathcal{O}_X are $\mathbf{0}$ -regular, we must have $\mathcal{V}_{\tau \cup \sigma} \cap \mathcal{R}^{d+d'} = \emptyset$. \square

The following example demonstrates that even when X is a smooth Fano toric variety the structure sheaf \mathcal{O}_X may not be $\mathbf{0}$ -regular.

Example 6.11. Let X be a blowup of \mathbb{P}^2 at three points. Specifically, the lattice points \mathcal{B} correspond to the columns of the matrix $\begin{bmatrix} 1 & 0 & -1 & -1 & 0 & 1 \\ 0 & 1 & 1 & 0 & -1 & -1 \end{bmatrix}$. The regular triangulation of \mathcal{B} is encoded in the irrelevant ideal

$$B = \langle x_1 x_2 x_3 x_4, x_2 x_3 x_4 x_5, x_3 x_4 x_5 x_6, x_1 x_4 x_5 x_6, x_1 x_2 x_5 x_6, x_1 x_2 x_3 x_6 \rangle.$$

We have $G = \mathbb{Z}^4$ and we may assume that \mathcal{A} is given by the columns of the matrix

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 1 & -1 \\ 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 & 1 \end{bmatrix}.$$

Let \mathcal{C} be the unique minimal generators of the monoid $\mathcal{H} = \mathcal{H}^{\text{sat}}$:

$$\mathcal{C} = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

It follows from Corollary 3.4 that

$$H^2(X, \mathcal{O}_X(-c_3 - c_5)) = H^2(X, \mathcal{O}_X(-a_1 - a_2 - a_3 - a_4 - a_5 - a_6)) = \tilde{H}^1(\Delta) \neq 0$$

which implies $\mathbf{0} \notin \text{reg}(\mathcal{O}_X)$.

The following remains an interesting open problem.

Problem 6.12. Let \mathcal{C} be a set of minimal generators of \mathcal{K} . Give a combinatorial characterization of all toric varieties X such that $\mathcal{K} \subseteq \text{reg}(\mathcal{O}_X)$.

We end this section with an upper semi-continuity result. This is well-known when $X = \mathbb{P}^d$. We write $\text{in}(I)$ for the initial ideal of an ideal $I \subseteq S$ with respect to some monomial order.

Proposition 6.13. *Let S be the homogeneous coordinate ring of a simplicial toric variety. If I is an ideal in S , then $\text{reg}(S/\text{in}(I)) \subseteq \text{reg}(S/I)$. Moreover, if I is B -saturated and $J = (\text{in}(I) : B^\infty)$ then $\text{reg}(S/J) \subseteq \text{reg}(S/I)$.*

Proof. Fix $j > 0$. If $\mathbf{x}^u f$ belongs to I for all $\mathbf{x}^u \in B^j$, then $\mathbf{x}^u \text{in}(f)$ belongs to $\text{in}(I)$. Hence, $f \in (I : B^\infty)$ implies that $\text{in}(f) \in (\text{in}(I) : B^\infty)$. We conclude that

$$\dim_{\mathbb{k}} H_B^0(S/I)_p \leq \dim_{\mathbb{k}} H_B^0(S/\text{in}(I))_p \quad \text{for all } p \in G.$$

[15], Theorem 15.17 gives a flat family over \mathbb{A}^1 whose general fiber is S/I and whose special fiber is $S/(\text{in}(I))$. Since $H_B^{i+1}(F) \cong \bigoplus_{p \in G} H^i(X, \mathcal{F}(p))$ for all $i \geq 1$, [23], Theorem 12.8 implies that $\dim_{\mathbb{k}} H_B^i(S/I)_p \leq \dim_{\mathbb{k}} H_B^i(S/\text{in}(I))_p$ for all $i > 1$. If \mathcal{I} denotes the sheaf of ideals corresponding to I , then the exact sequence (6.3.1) gives

$$\dim_{\mathbb{k}} H_B^0(S/I)_p + \dim_{\mathbb{k}} H^0(X, (\mathcal{O}_X/\mathcal{I})(p)) = \dim_{\mathbb{k}}(S/I)_p + \dim_{\mathbb{k}} H_B^1(S/I)_p.$$

Since the terms on left-hand side do not decrease and $\dim_{\mathbb{k}}(S/I)_p$ is constant when passing to an initial ideal, we see that $\dim_{\mathbb{k}} H_B^1(S/I)_p$ also does not decrease when passing to an initial ideal. Thus, we conclude that $H_B^i(S/\text{in}(I))_p = 0$ implies $H_B^i(S/I)_p = 0$ and the first assertion follows. Since $H_B^0(S/J) = 0$ and $H_B^i(S/J) = H_B^i(S/\text{in}(I))$ for $i > 0$, this also establishes the second assertion. \square

7. Syzygies and chain complexes

The final section of the paper examines the relationship between the regularity of M and the syzygies of M . We give a combinatorial formula, involving only the degrees of the syzygies and the regularity of S , for a subset of $\text{reg}(M)$. When S is the homogeneous coor-

dinate ring of weighted projective space, this subset actually equals $\text{reg}(M)$. In other words, we recover the characterization of regularity in terms of Betti numbers in this case; see [3], Theorem 5.5. We extend this formula to coherent \mathcal{O}_X -modules. Changing directions, we then describe chain complexes associated to certain elements of $\text{reg}(M)$. At the level of \mathcal{O}_X -modules, we obtain a locally free resolution of \mathcal{F} from specific elements in $\text{reg}(\mathcal{F})$.

If \mathbf{E} is a free resolution of the module M , then there is a spectral sequence that computes the local cohomology of M from \mathbf{E} , namely $E_2^{i,j} := H^i(H_B^j(\mathbf{E})) \Rightarrow H_B^{i+j}(M)$. When S is the homogeneous coordinate ring of weighted projective space, this spectral sequence degenerates because $H_B^i(S)$ vanishes unless $i = d + 1$; see Remark 3.3. As a consequence, there is a simple characterization of the regularity of M in terms of the degrees of the syzygies. In contrast, Remark 3.3 also shows that nonvanishing local cohomology of S is not typically concentrated in a single cohomological degree. Hence, the spectral sequence does not degenerate and one cannot expect as simple a relationship between regularity and syzygies in the general situation. Despite this, the syzygies of M do provide a method for approximating the regularity of M which captures the description in the special case. Moreover, this technique also works on a larger class of chain complexes associated to M .

We start by describing how regularity behaves in short exact sequences.

Lemma 7.1. *If $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is a short exact sequence of finitely generated G -graded S -modules, then we have the following:*

- (1) $\text{reg}^i(M') \cap \text{reg}^i(M'') \subseteq \text{reg}^i(M)$,
- (2) $\left(\bigcup_{1 \leq j \leq \ell} (-\mathbf{c}_j + \text{reg}^{i+1}(M')) \right) \cap \text{reg}^i(M) \subseteq \text{reg}^i(M'')$ and
- (3) $\text{reg}^i(M) \cap \left(\bigcap_{1 \leq j \leq \ell} (\mathbf{c}_j + \text{reg}^{i-1}(M'')) \right) \subseteq \text{reg}^i(M')$.

Proof. The associated long exact sequence in cohomology contains the exact sequence

$$(7.1.1) \quad H_B^{i-1}(M'')_{\mathbf{p}} \rightarrow H_B^i(M')_{\mathbf{p}} \rightarrow H_B^i(M)_{\mathbf{p}} \rightarrow H_B^i(M'')_{\mathbf{p}} \rightarrow H_B^{i+1}(M')_{\mathbf{p}}.$$

Suppose that both M' and M'' are \mathbf{m} -regular from level k . This means that the second and fourth modules in (7.1.1) vanish for all $i \geq k$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$. Hence, the third module in (7.1.1) also vanishes for all $i \geq k$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$ which implies that M is \mathbf{m} -regular from level k .

Similarly, if M is \mathbf{m} -regular from level k and M' is $(\mathbf{m} + \mathbf{c}_j)$ -regular from level $(k + 1)$ for some $1 \leq j \leq \ell$, then the third and fifth modules in (7.1.1) vanish for all $i \geq k$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$. It follows that the fourth module in (7.1.1) also vanishes under the same conditions. Thus, M'' is \mathbf{m} -regular from level k .

Finally, if M is \mathbf{m} -regular from level k and M'' is $(\mathbf{m} - \mathbf{c}_j)$ -regular from level $(k - 1)$ for every $1 \leq j \leq \ell$, then the first and third modules in (7.1.1) vanish for all $i \geq 0$ and all $\mathbf{p} \in \mathbf{m} + \mathbb{N}\mathcal{C}[1 - i]$. Hence, the second module in (7.1.1) vanishes under the same conditions and therefore $\mathbf{m} \in \text{reg}^k(M')$. \square

The difference between the union in Part (2) and the intersection in Part (3) introduces an asymmetry in working with short exact sequences. In many cases, this prevents one from giving a simple characterization of \mathbf{m} -regularity in terms of minimal free resolutions. However, when \mathcal{C} consists of a single element, as in the standard graded case, this obstruction is not present. In this case, we have the more symmetric:

$$(-\mathbf{c}_1 + \text{reg}^{i+1}(M')) \cap \text{reg}^i(M) \subseteq \text{reg}^i(M)$$

and $(\mathbf{c}_1 + \text{reg}^{i-1}(M'')) \cap \text{reg}^i(M) \subseteq \text{reg}^i(M')$.

The approach used in the proof of Lemma 7.1 also leads to an analogous result for a short exact sequence of coherent \mathcal{O}_X -modules. However, there is a notable change in Part (3). Specifically, we must assume that $i > 1$ because the hypothesis that \mathcal{F} is \mathbf{m} -regular does not place any conditions on $H^0(X, \mathcal{F}(\mathbf{p}))$.

The next theorem provides a method for estimating the regularity of a module M from certain chain complexes associated to M . We say a chain complex of S -modules has *B-torsion homology* if every homology module is a B -torsion module.

Theorem 7.2. *Let $\mathbf{E} := \cdots \rightarrow E_3 \xrightarrow{\partial_3} E_2 \xrightarrow{\partial_2} E_1 \xrightarrow{\partial_1} E_0 \rightarrow 0$ be a chain complex of finitely generated G -graded S -modules with B -torsion homology. If $\partial_0 : E_0 \rightarrow M$ is a surjective map then we have*

$$\bigcup_{\phi: [d+1] \rightarrow [\ell]} \left(\bigcap_{0 \leq i \leq d+1} (-\mathbf{c}_{\phi(1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(E_i)) \right) \subseteq \text{reg}(M),$$

where the union is over all functions $\phi : [d + 1] \rightarrow [\ell]$.

Proof. Fix a function $\phi : [d + 1] \rightarrow [\ell]$. We claim that for all $k \geq 0$

$$\bigcap_{k \leq i \leq d+1} (-\mathbf{c}_{\phi(k+1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(E_i)) \subseteq \text{reg}^k(\text{Im } \partial_k).$$

Since $\text{Im } \partial_0 = M$, this will prove the theorem. We establish the claim by using a descending induction on k . Since $H_B^j(\text{Im } \partial_k)$ vanishes for all $j > d + 1$, we have $\text{reg}^k(\text{Im } \partial_k) = G$ for $k > d + 1$ and the claim holds. Suppose $k \leq d + 1$ and consider the exact sequence $0 \rightarrow H_k(\mathbf{E}) \rightarrow E_k/\text{Im } \partial_{k+1} \rightarrow \text{Im } \partial_k \rightarrow 0$. Because $H_k(\mathbf{E})$ is a B -torsion module, the associated long exact sequence in cohomology implies that $H_B^j(E_k/\text{Im } \partial_{k+1}) = H_B^j(\text{Im } \partial_k)$ for all $j > 0$ and that $H_B^0(E_k/\text{Im } \partial_{k+1})$ surjects onto $H_B^0(\text{Im } \partial_k)$. It follows that

$$\text{reg}^k(E_k/\text{Im } \partial_{k+1}) \subseteq \text{reg}^k(\text{Im } \partial_k) \quad \text{for all } k \geq 0.$$

Applying Lemma 7.1 to the short exact sequence $0 \rightarrow \text{Im } \partial_{k+1} \rightarrow E_k \rightarrow E_k/\text{Im } \partial_{k+1} \rightarrow 0$ gives $\text{reg}^{k+1}(\text{Im } \partial_{k+1})[-1] \cap \text{reg}^k(E_k) \subseteq \text{reg}^k(E_k/\text{Im } \partial_{k+1})$. Since the induction hypothesis implies that $\bigcap_{k+1 \leq i \leq d+1} (-\mathbf{c}_{\phi(k+2)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(E_i)) \subseteq \text{reg}^{k+1}(\text{Im } \partial_{k+1})$, we have

$\bigcap_{k+1 \leq i \leq d+1} (-\mathbf{c}_{\phi(k+1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(E_i)) \subseteq \text{reg}^{k+1}(\text{Im } \partial_{k+1})[-1]$. Combining these three inclusions, we obtain $\bigcap_{k \leq i \leq d+1} (-\mathbf{c}_{\phi(k+1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(E_i)) \subseteq \text{reg}^k(\text{Im } \partial_k)$. \square

Since the proof of Theorem 7.2 only used Part (2) of Lemma 7.1, the same approach leads to a formula for estimating the regularity of a sheaf \mathcal{F} from a resolution. We leave the details to the interested reader. Because it is frequently used in applications, we explicitly state the following special case of Theorem 7.2.

Corollary 7.3. *If $0 \rightarrow E_s \rightarrow \cdots \rightarrow E_2 \rightarrow E_1 \rightarrow E_0$ is a free resolution of the module M with $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{q}_{i,j})$ for some $\mathbf{q}_{i,j} \in G$, then we have*

$$(7.3.2) \quad \bigcup_{\phi: [d+1] \rightarrow [l]} \left(\bigcap_{\substack{0 \leq i \leq \min\{d+1, s\} \\ 1 \leq j \leq h_i}} (\mathbf{q}_{i,j} - \mathbf{c}_{\phi(1)} - \cdots - \mathbf{c}_{\phi(i)} + \text{reg}^i(S)) \right) \subseteq \text{reg}(M).$$

Proof. Since $\text{reg}^i(E_i) = \bigcap_{1 \leq j \leq h_i} (\mathbf{q}_{i,j} + \text{reg}^i(S))$, (7.3.2) follows from Theorem 7.2. \square

The next three examples illustrate this corollary.

Example 7.4. Suppose S is the homogeneous coordinate ring of weighted projective space; see Example 2.1. Let $\mathcal{C} = \{\mathbf{1}\}$ and let $0 \rightarrow E_s \rightarrow \cdots \rightarrow E_2 \rightarrow E_1 \rightarrow E_0$ be the minimal free resolution of the module M where $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{q}_{i,j})$ for some $\mathbf{q}_{i,j} \in \mathbb{Z}$. Hilbert’s syzygy theorem implies that $s \leq n = d + 1$. Hence, Example 4.2 and (7.3.2) imply that $\max_{i,j} \{\mathbf{q}_{i,j} - i + n - \mathbf{a}_1 - \cdots - \mathbf{a}_n\} \in \text{reg}(M)$. In particular, if S has the standard grading, then M is $(\max_{i,j} \{\mathbf{q}_{i,j} - i\})$ -regular.

Example 7.5. Let S be the homogeneous coordinate ring of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. This means that \mathcal{B} corresponds to the columns of the matrix

$$\begin{bmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix},$$

and the irrelevant ideal is $B = \langle x_1, x_2 \rangle \cap \langle x_3, x_4 \rangle \cap \langle x_5, x_6 \rangle$. Hence, $G = \mathbb{Z}^3$ and we may assume that \mathcal{A} is given by the columns of the matrix

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}.$$

The polynomial ring $S = \mathbb{k}[x_1, \dots, x_6]$ has the corresponding \mathbb{Z}^3 -grading and $\mathcal{H} = \mathcal{H}^{\text{sat}} = \mathbb{N}^3$. If $\mathcal{C} = \{\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_3\} \subset \mathcal{H}$ where \mathbf{c}_i is the i th standard basis vector in \mathbb{Z}^3 , then Corollary 3.4 implies that $\text{reg}(S) = \mathbb{N}^3$.

Consider the module $M = S/I$ where $I = \langle x_1 - x_2, x_3 - x_4, x_5 - x_6 \rangle$. Since I is a prime ideal of codimension 3, the subvariety of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ defined by I is a single point. The minimal free resolution of M has the form

$$0 \rightarrow S(-\mathbf{c}_1 - \mathbf{c}_2 - \mathbf{c}_3) \rightarrow \bigoplus_{1 \leq i < j \leq 3} S(-\mathbf{c}_i - \mathbf{c}_j) \rightarrow \bigoplus_{1 \leq i \leq 3} S(-\mathbf{c}_i) \rightarrow S.$$

A calculation using Corollary 7.3 shows that $\bigcup_{1 \leq i < j \leq 3} (\mathbf{e}_i + \mathbf{e}_j + \mathbb{N}^3) \subseteq \text{reg}(M)$. Note that this is not the entire regularity of M . Proposition 6.7 shows that $\text{reg}(M) = \mathbb{N}^3$.

Example 7.6. As in Example 7.5, we assume that S is the homogeneous coordinate ring of the toric variety $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and we choose \mathcal{C} to be the set of standard basis vectors in \mathbb{Z}^3 . Consider $M = S/I$ where

$$I = \langle x_1 - x_2, x_3 - x_4, x_5 - x_6 \rangle \cap \langle x_1 - 2x_2, x_3 - 2x_4, x_5 - 2x_6 \rangle.$$

The B -saturated ideal I corresponds to two distinct points on $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$. The minimal free resolution of M has the form

$$0 \rightarrow \bigoplus_{\mathbf{q} \in \mathcal{D}_5} S(-\mathbf{q}) \rightarrow \bigoplus_{\mathbf{q} \in \mathcal{D}_4} S(-\mathbf{q}) \rightarrow \bigoplus_{\mathbf{q} \in \mathcal{D}_3} S(-\mathbf{q}) \rightarrow \bigoplus_{\mathbf{q} \in \mathcal{D}_2} S(-\mathbf{q}) \rightarrow \bigoplus_{\mathbf{q} \in \mathcal{D}_1} S(-\mathbf{q}) \rightarrow S$$

where, for $1 \leq i \leq 5$, the set \mathcal{D}_i is given by the column vectors of the matrix D_i and

$$D_1 = \begin{bmatrix} 2 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 2 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 2 \end{bmatrix},$$

$$D_2 = \begin{bmatrix} 2 & 2 & 2 & 2 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 2 & 2 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 2 & 2 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 2 & 1 & 1 & 2 & 2 \end{bmatrix},$$

$$D_3 = \begin{bmatrix} 2 & 2 & 2 & 2 & 2 & 2 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 & 1 & 0 & 2 & 2 & 2 & 2 & 1 & 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 1 & 1 & 2 & 1 & 1 & 1 & 1 & 2 & 2 & 2 & 2 & 2 \end{bmatrix},$$

$$D_4 = \begin{bmatrix} 2 & 2 & 2 & 2 & 1 & 1 \\ 2 & 2 & 1 & 1 & 2 & 2 \\ 1 & 1 & 2 & 2 & 2 & 2 \end{bmatrix}, \quad D_5 = \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}.$$

Observe that the length of the minimal free resolution is greater than $d + 1$. Applying Corollary 7.3, we deduce that

$$\left(\begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix} + \mathbb{N}^3 \right) \cup \left(\begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix} + \mathbb{N}^3 \right) \cup \left(\begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} + \mathbb{N}^3 \right) \subset \text{reg}(M).$$

However, Proposition 6.7 shows that

$$\text{reg}(M) = \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + \mathbb{N}^3 \right) \cup \left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + \mathbb{N}^3 \right) \cup \left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \mathbb{N}^3 \right).$$

To describe a partial converse for Theorem 7.2, we concentrate on the situation arising from smooth toric varieties. Specifically, we assume that $\mathcal{K} = \mathcal{K}^{\text{sat}}$ and that \mathcal{C} is a finite generating set for the monoid \mathcal{K} .

Lemma 7.7. *Let $\mathbb{N}\mathcal{C} = \mathcal{K} = \mathcal{K}^{\text{sat}}$. If M is \mathbf{m} -regular then $M|_{(\mathbf{m}+\mathcal{K})}$ is also \mathbf{m} -regular.*

Proof. Suppose that M is \mathbf{m} -regular and let M' be the quotient module $M/M|_{(\mathbf{m}+\mathcal{K})}$. Since $\mathbb{N}\mathcal{C} = \mathcal{K} = \mathcal{K}^{\text{sat}}$, Lemma 6.8 establishes that M' is a B -torsion module. Hence, the long exact sequence associated to (6.8.2) implies that $H_B^i(M|_{(\mathbf{m}+\mathcal{K})}) = H_B^i(M)$ for all $i > 1$ and $H_B^1(M|_{(\mathbf{m}+\mathcal{K})})_p = H_B^1(M)_p$ for all $p \in \mathbf{m} + \mathcal{K}$. For $k > 0$, we deduce that M is \mathbf{m} -regular from level k if and only if the submodule $M|_{(\mathbf{m}+\mathcal{K})}$ is \mathbf{m} -regular from level k . The long exact sequence also shows that $H_B^0(M|_{(\mathbf{m}+\mathcal{K})})$ is a submodule of $H_B^0(M)$. Thus, if $H_B^0(M)_p = 0$ then $H_B^0(M|_{(\mathbf{m}+\mathcal{K})})_p$ also vanishes. Therefore, $M|_{(\mathbf{m}+\mathcal{K})}$ is \mathbf{m} -regular. \square

Theorem 7.8. *Assume that $\mathbb{N}\mathcal{C} = \mathcal{K} = \mathcal{K}^{\text{sat}}$. If $\mathbf{c} \in \text{reg}(S) \cap \left(\bigcap_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathcal{K}) \right)$ and $\mathbf{m} \in \text{reg}(M)$ then there exists*

(1) *a chain complex $\cdots \rightarrow E_3 \xrightarrow{\partial_3} E_2 \xrightarrow{\partial_2} E_1 \xrightarrow{\partial_1} E_0$ with B -torsion homology and $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{m} - i\mathbf{c})$, and*

(2) *a surjective map $\partial_0 : E_0 \rightarrow M|_{(\mathbf{m}+\mathbb{N}\mathcal{C})}$.*

Proof. We prove by induction on k that there exists a chain complex with B -torsion homology

$$(7.8.3) \quad 0 \longrightarrow M_k \longrightarrow E_{k-1} \xrightarrow{\partial_{k-1}} \cdots \xrightarrow{\partial_1} E_0 \xrightarrow{\partial_0} M \xrightarrow{\partial_{-1}} 0$$

such that $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{m} - i\mathbf{c})$, $M_k = \text{Ker } \partial_{k-1}$ and $\mathbf{m} + k\mathbf{c} \in \text{reg}(M_k)$. Since $M = M_0$ and M is \mathbf{m} -regular, the first step in the induction holds.

Suppose $k \geq 0$. Since $\mathbb{N}\mathcal{C} = \mathcal{K} = \mathcal{K}^{\text{sat}}$, Lemma 7.7 and Lemma 6.8 show that $\text{reg}(M_k) \subseteq \text{reg}(M_k|_{(\mathbf{m}+k\mathbf{c}+\mathbb{N}\mathcal{C})})$ and that $M_k/M_k|_{(\mathbf{m}+k\mathbf{c}+\mathbb{N}\mathcal{C})}$ is a B -torsion module. The induction hypothesis states that M_k is $(\mathbf{m} + k\mathbf{c})$ -regular and Theorem 5.4 implies that $M_k|_{(\mathbf{m}+k\mathbf{c}+\mathbb{N}\mathcal{C})}$ is generated in degree $\mathbf{m} + k\mathbf{c}$. It follows that there exists a surjective map $\partial_k : E_k \rightarrow M_k|_{(\mathbf{m}+k\mathbf{c}+\mathbb{N}\mathcal{C})}$ where $E_k := \bigoplus_{1 \leq j \leq b_k} S(-\mathbf{m} - k\mathbf{c})$. Setting $M_{k+1} := \text{Ker } \partial_k$, we obtain the short exact sequence

$$(7.8.4) \quad 0 \rightarrow M_{k+1} \rightarrow E_k \rightarrow M_k|_{(\mathbf{m}+k\mathbf{c}+\mathbb{N}\mathcal{C})} \rightarrow 0.$$

Combining (7.8.3) and (7.8.4) gives the chain complex with B -torsion homology

$$0 \rightarrow M_{k+1} \rightarrow E_k \xrightarrow{\partial_k} \cdots \xrightarrow{\partial_1} E_0 \xrightarrow{\partial_0} M \xrightarrow{\partial_{-1}} 0$$

where $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{m} - i\mathbf{c})$ and $M_{k+1} = \text{Ker } \partial_k$. It remains to show that M_{k+1} is $(\mathbf{m} + (k+1)\mathbf{c})$ -regular. Applying Lemma 7.1 to (7.8.4) yields

$$(\mathbf{m} + k\mathbf{c} + \operatorname{reg}(S)) \cap \bigcap_{1 \leq i \leq \ell} (\mathbf{c}_i + \operatorname{reg}(M_k)) \subseteq \operatorname{reg}(M_{k+1}).$$

Our choice of \mathbf{c} guarantees that $\mathbf{m} + (k+1)\mathbf{c}$ lies in both $\bigcap_{1 \leq i \leq \ell} (\mathbf{c}_i + \operatorname{reg}(M_k))$ and $\mathbf{m} + k\mathbf{c} + \operatorname{reg}(S)$ which completes the proof. \square

Proof of Theorem 1.5. In the introduction, X was smooth so $\mathcal{K} = \mathcal{K}^{\text{sat}}$. We also assumed that \mathcal{C} was the set of minimal generators for \mathcal{K} . Hence, Theorem 7.2 establishes Part (1) and Theorem 7.8 proves Part (2). \square

This theorem leads to a “linear” resolution of \mathcal{O}_X -modules.

Corollary 7.9. *Assume that $\mathbb{N}\mathcal{C} = \mathcal{K} = \mathcal{K}^{\text{sat}}$. If $\mathbf{c} \in \operatorname{reg}(\mathcal{O}_X) \cap \left(\bigcap_{1 \leq j \leq \ell} (\mathbf{c}_j + \mathcal{K}) \right)$ and $\mathbf{m} \in \operatorname{reg}(\mathcal{F})$ then there is an exact sequence*

$$\cdots \rightarrow \mathcal{E}_3 \rightarrow \mathcal{E}_2 \rightarrow \mathcal{E}_1 \rightarrow \mathcal{E}_0 \rightarrow \mathcal{F} \rightarrow 0$$

where $\mathcal{E}_i = \bigoplus_{1 \leq j \leq h_i} \mathcal{O}_X(-\mathbf{m} - i\mathbf{c}_j)$.

Proof. Consider the G -graded S -module

$$M = \left(\bigoplus_{p \in G} H^0(X, \mathcal{F}(p)) \right) \Big|_{(\mathbf{m} + \mathcal{K})}.$$

Lemma 6.8 implies that $\tilde{M} = \mathcal{F}$ and Proposition 6.4 shows that M is \mathbf{m} -regular. Theorem 7.8 produces a chain complex $\cdots \rightarrow E_3 \rightarrow E_2 \rightarrow E_1 \rightarrow E_0 \rightarrow M \rightarrow 0$ with B -torsion homology and $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{m} - i\mathbf{c}_j)$. Applying the functor $F \mapsto \tilde{F}$, we obtain a chain complex of the desired form. Moreover, Proposition 6.1 implies that the homology of this complex consists of the zero sheaf. Therefore, the chain complex is a locally free resolution of \mathcal{F} . \square

We illustrate Theorem 7.8 with the following examples. The first example shows that Theorem 7.8 is a converse to Theorem 7.2 in the standard graded case.

Example 7.10. Let S have the standard grading, let $B = \langle x_1, \dots, x_n \rangle$, and let $\mathcal{C} = \{\mathbf{1}\}$. Since $\operatorname{reg}(S) = \mathbb{N}$, we have $\mathbf{c} = \mathbf{1} \in \operatorname{reg}(S) \cap (\mathbf{1} + \mathcal{K})$. Hence, if $\mathbf{m} \in \operatorname{reg}(M)$, then Theorem 7.8 implies that there is a chain complex $\mathbf{E} := \cdots \rightarrow E_2 \rightarrow E_1 \rightarrow E_0$ with B -torsion homology and $E_i = \bigoplus_{1 \leq j \leq h_i} S(-\mathbf{m} - i)$ and there is a surjective map $E_0 \rightarrow M|_{(\mathbf{m} + \mathbb{N})}$. Conversely, given such a chain complex, Example 7.4 shows that $\mathbf{m} \in \operatorname{reg}(M)$. In fact, [4], Exercise 1.4.24 proves that \mathbf{E} must be a resolution.

Example 7.11. Let M be the module in Example 7.5 and let

$$\mathbf{c} := \mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3 \in \operatorname{reg}(S) \cap \left(\bigcap_{1 \leq j \leq 3} (\mathbf{c}_j + \mathbb{N}\mathcal{C}) \right).$$

If $\mathbf{m} := \mathbf{0} \in \operatorname{reg}(M)$, then Theorem 7.8 produces a chain complex with B -torsion homology of the form

$$\cdots \rightarrow \bigoplus_i S(-3\mathbf{c}) \rightarrow \bigoplus_i S(-2\mathbf{c}) \rightarrow \bigoplus_{1 \leq i \leq 7} S(-\mathbf{c}) \rightarrow S.$$

However, applying Theorem 7.2 to this chain complex yields

$$\bigcup_{\substack{p_1, p_2, p_3 \in \mathbb{N} \\ p_1 + p_2 + p_3 = 8}} \left(\begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix} + \mathbb{N}^3 \right) \subset \text{reg}(M),$$

and the smaller set does not contain $\mathbf{0}$.

References

- [1] *D. Bayer and M. Stillman*, A criterion for detecting m -regularity, *Invent. Math.* **87** (1987), 1–11.
- [2] *D. Bayer and B. Sturmfels*, Cellular resolutions of monomial modules, *J. reine angew. Math.* **502** (1998), 123–140.
- [3] *D. Benson*, Dickson invariants, regularity and computation in group cohomology, arXiv:math.GR/0303187.
- [4] *L. J. Billera, I. M. Gel'fand, and B. Sturmfels*, Duality and minors of secondary polyhedra, *J. Combin. Th. (B)* **57** (1993), 258–268.
- [5] *M. P. Brodmann and R. Y. Sharp*, Local cohomology: an algebraic introduction with geometric applications, *Cambridge Stud. Adv. Math.* **60**, Cambridge Univ. Press, Cambridge 1998.
- [6] *W. Bruns, J. Gubeladze, and N. V. Trung*, Normal polytopes, triangulations, and Koszul algebras, *J. reine angew. Math.* **485** (1997), 123–160.
- [7] *W. Bruns and J. Herzog*, Cohen-Macaulay rings, *Cambridge Stud. Adv. Math.* **39**, Cambridge Univ. Press, Cambridge 1993.
- [8] *W. Bruns and R. Koch*, Normaliz: Computing the normalization of affine semigroups, <ftp://ftp.mathematik.uni-osnabrueck.de/pub/osm/kommalg/software/>.
- [9] *J. V. Chipalkatti*, A generalization of Castelnuovo regularity to Grassmann varieties, *Manuscr. Math.* **102** (2000), 447–464.
- [10] *D. A. Cox*, Equations of Parametric Surfaces via Syzygies, arXiv:math.AG/0010107.
- [11] *D. A. Cox*, The homogeneous coordinate ring of a toric variety, *J. Alg. Geom.* **4** (1995), 17–50.
- [12] *D. A. Cox*, Recent developments in toric geometry, Algebraic geometry—Santa Cruz 1995, *Proc. Sympos. Pure Math.* **62**, Amer. Math. Soc., Providence, RI (1997), 389–436.
- [13] *L. Ein and R. Lazarsfeld*, Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension, *Invent. Math.* **111** (1993), 51–67.
- [14] *D. Eisenbud and S. Goto*, Linear free resolutions and minimal multiplicity, *J. Algebra* **88** (1984), 89–133.
- [15] *D. Eisenbud*, Commutative algebra with a view toward algebraic geometry, *Grad. Texts Math.* **150**, Springer-Verlag, New York 1995.
- [16] *D. Eisenbud, M. Mustață, and M. Stillman*, Cohomology on toric varieties and local cohomology with monomial supports, *J. Symb. Comput.* **29** (2000), 583–600.
- [17] *W. Fulton*, Introduction to toric varieties, *Ann. Math. Stud.* **131**, Princeton Univ. Press, Princeton, NJ, 1993.
- [18] *G. Gotzmann*, Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes, *Math. Z.* **158** (1978), 61–70.
- [19] *D. R. Grayson and M. E. Stillman*, Macaulay 2, a software system for research in algebraic geometry, <http://www.math.uiuc.edu/Macaulay2>.
- [20] *L. Gruson, R. Lazarsfeld, and C. Peskine*, On a theorem of Castelnuovo, and the equations defining space curves, *Invent. Math.* **72** (1983), 491–506.
- [21] *H. T. Hà and A. Van Tuyl*, The regularity of points in multi-projective spaces, math.AC/0211263.
- [22] *M. Haiman and B. Sturmfels*, Multigraded Hilbert schemes, arXiv:math.AG/0201271.
- [23] *R. Hartshorne*, Algebraic geometry, *Grad. Texts Math.* **52**, Springer-Verlag, New York 1977.
- [24] *T. Hausel and B. Sturmfels*, Toric Hyperkähler Varieties, *Doc. Math.* **7** (2002), 495–534.
- [25] *J. W. Hoffman and H. H. Wang*, Castelnuovo-Mumford Regularity in Biprojective Spaces, arXiv:math.AG/0212033.
- [26] *A. Iarrobino and S. L. Kleiman*, Appendix C, in: A. Iarrobino and V. Kanev, Power sums, Gorenstein algebras, and determinantal loci, *Lect. Notes Math.* **1721**, Springer-Verlag, Berlin 1999.

- [27] *R. Lazarsfeld*, Positivity in algebraic geometry, to appear.
- [28] *D. Maclagan* and *G. G. Smith*, Uniform bounds on multigraded regularity, arXiv:math.AG/0305215.
- [29] *E. Miller*, The Alexander duality functors and local duality with monomial support, *J. Algebra* **231** (2000), 180–234.
- [30] *D. Mumford*, Lectures on curves on an algebraic surface, Princeton Univ. Press, Princeton, N.J., 1966.
- [31] *J. R. Munkres*, Elements of algebraic topology, Addison-Wesley Publishing Company, Menlo Park, CA, 1984.
- [32] *T. Oda*, Open problems on Minkowski sums of polytopes, 1997, Tôhoku University preprint.
- [33] *G. Pareschi* and *M. Popa*, Regularity on abelian varieties I, *J. Amer. Math. Soc.* **16** (2003), 285–302.
- [34] *V. Reiner*, *V. Welker*, and *K. Yanagawa*, Local cohomology of Stanley-Reisner rings with supports in general monomial ideals, *J. Algebra* **244** (2001), 706–736.
- [35] *P. C. Roberts*, Multiplicities and Chern classes in local algebra, Cambridge Tracts Math. **133**, Cambridge Univ. Press, Cambridge 1998.
- [36] *G. M. Ziegler*, Lectures on polytopes, Grad. Texts Math. **152**, Springer-Verlag, New York 1995.

Department of Mathematics, Stanford University, Stanford, CA 94305, USA
e-mail: maclagan@math.stanford.edu

Department of Mathematics, Barnard College, Columbia University, New York, NY 10027, USA
e-mail: ggsmith@math.columbia.edu

Eingegangen 15. Mai 2003, in revidierter Fassung 30. Juni 2003