

1. Recall that an ideal $I \subseteq k[Z_0, \dots, Z_n]$ is called a *homogeneous ideal* if, for every $f \in I$, when we write out f as a sum of homogeneous pieces, $f = F_0 + F_1 + F_2 + \dots + F_d$, then each F_j is also in I .

In this problem we will show that an ideal I is homogeneous if and only if I can be generated by homogeneous polynomials, i.e., if and only if there are homogeneous polynomials $G_1, \dots, G_s \in k[Z_0, \dots, Z_n]$ such that $I = \langle G_1, \dots, G_s \rangle$.

- (a) Assume that I is a homogeneous ideal. Show that I may be generated by homogeneous polynomials. (SUGGESTION: since I is an ideal in $k[Z_0, \dots, Z_n]$ it may be generated by finitely many polynomials. Apply the ‘homogeneous ideal’ condition to these generators.)
- (b) Suppose that $I = \langle G_1, \dots, G_s \rangle$ with each G_i a homogeneous polynomial. Show that I is a homogeneous ideal. (SUGGESTION: Any $f \in I$ can be written as $f = h_1 G_1 + \dots + h_s G_s$ with the h_i polynomials in Z_0, \dots, Z_n . Write each h_i as a sum of homogeneous pieces, expand the sum $\sum h_i G_i$, collect pieces of the same degree and compare with the homogeneous pieces of f .)
- (c) Is the ideal $\langle X^3 - 5XZ^2 + Y^2 + XY, X^3 - 5XZ^2 - Y^2 - XY \rangle \subset \mathbb{Q}[X, Y, Z]$ homogeneous?

2. Recall that a subvariety $Y \subseteq \mathbb{A}^n$ is called a *cone* if whenever $p \in Y$ then $\lambda p \in Y$ for all $\lambda \in k^*$, where λp means the point obtained by scaling all the coordinates of p by λ . In this problem we will show that Y is a cone if and only if J_Y , the ideal of Y , is a homogeneous ideal. For this question we assume that k is an infinite field.

First suppose that Y is a cone, let f be an element of J_Y and write f as a sum of homogeneous pieces, $f = F_0 + F_1 + \dots + F_d$, with each F_j homogeneous of degree j . To show that J_Y is a homogeneous ideal, we need to show that each $F_j \in J_Y$.

- (a) Explain why it is sufficient to show that $F_j(p) = 0$ for all $p \in Y$.
- (b) Fix $p \in Y$. By considering $f(\lambda p)$, explain why

$$0 = F_0(p) + \lambda F_1(p) + \lambda^2 F_2(p) + \dots + \lambda^d F_d(p)$$

for all $\lambda \in k^*$.

- (c) Considering the expression in (b) as a polynomial in λ , explain why we must have $F_j(p) = 0$ for each j , and hence (from the reductions above) that J_Y is a homogeneous ideal.

- (d) Now prove the other direction : assume that $Y \subseteq \mathbb{A}^n$ is a variety such that J_Y is a homogeneous ideal, and prove that Y is a cone. (The equivalence in question 1 may help.)

3. Suppose that $U_0, U_1,$ and U_2 are the standard open subsets in \mathbb{P}^2 , and that we have varieties $Y_0 \subset U_0, Y_1 \subset U_1,$ and $Y_2 \subset U_2,$ which agree on intersections. This means that $Y_i|_{U_i \cap U_j} = Y_j|_{U_i \cap U_j}$ for any i, j . In this question we will prove that there is a homogeneous ideal I in $k[X, Y, Z]$ so that if we set $Y = V(I)$ then $Y \cap U_i = Y_i$ for $i = 0, 1, 2$. I.e., we will show that if we define a subvariety of \mathbb{P}^2 as something obtained by glueing together affine varieties on the pieces, then this agrees with our definition of subvariety as something obtained by homogeneous polynomials.

Since $Y_0, Y_1,$ and Y_2 are each affine varieties (in $U_0, U_1,$ and U_2 respectively) each of them are given by ideals in their respective polynomial rings. Let $I_0, I_1,$ and I_2 be these ideals. Then let $\tilde{I}_0, \tilde{I}_1,$ and \tilde{I}_2 be the homogenization of these ideals; namely the ideal obtained by homogenizing the polynomials in $I_0, I_1,$ and I_2 respectively. These ideals have the property that $V(\tilde{I}_j) \cap U_j = Y_j$ for each j . In other words, they each define projective varieties which restrict (separately) to the varieties we want on one of the open sets. However we do not know that $V(\tilde{I}_j) \cap U_i = Y_i$ when $i \neq j$, so these ideals by themselves do not solve the problem.

Recall that $U_0 = \mathbb{P}^2 \setminus \{X = 0\}, U_1 = \mathbb{P}^2 \setminus \{Y = 0\},$ and $U_2 = \mathbb{P}^2 \setminus \{Z = 0\}$. Let $\tilde{I}_0 X$ be the ideal obtained by multiplying all the elements of \tilde{I}_0 by X . If we look at $V(\tilde{I}_0 X)$, this variety contains all the points of $X = 0$ (i.e., all the points off of U_0), while we still have $V(\tilde{I}_0 X) \cap U_0 = Y_0$. Similar statements hold for $V(\tilde{I}_1 Y)$ and $V(\tilde{I}_2 Z)$ (with similar definitions for $\tilde{I}_1 Y$ and $\tilde{I}_2 Z$).

Finally define $I = \tilde{I}_0 X + \tilde{I}_1 Y + \tilde{I}_2 Z$, and set $Y = V(I)$. By our relation between subvarieties and geometric operations, this means that

$$Y = V(I) = V(\tilde{I}_0 X) \cap V(\tilde{I}_1 Y) \cap V(\tilde{I}_2 Z).$$

We now want to show that $Y \cap U_i = Y_i$ for each i . By symmetry of the construction it is enough to do this for $i = 0$.

(a) Show that $Y \cap U_0 \subseteq Y_0$.

(b) Show that $Y_0 \subseteq Y \cap U_0$.

The proofs of both statements involve only elementary considerations about intersections, and inclusions, and the way that Y was defined. In particular, part (a) should be very straightforward. For part (b) you will need the condition that $Y_i|_{U_i \cap U_j} = Y_j|_{U_i \cap U_j}$.