


1. If f is a function on \mathbb{R}^3 , and $\mathbf{F} = (F_1, F_2, F_3)$ a vector field on \mathbb{R}^3 , then $f\mathbf{F} = (fF_1, fF_2, fF_3)$, so

$$\begin{aligned} \operatorname{Div}(f\mathbf{F}) &= \frac{\partial}{\partial x}(fF_1) + \frac{\partial}{\partial y}(fF_2) + \frac{\partial}{\partial z}(fF_3) \\ &= \left(\frac{\partial f}{\partial x}F_1 + f\frac{\partial F_1}{\partial x} \right) + \left(\frac{\partial f}{\partial y}F_2 + f\frac{\partial F_2}{\partial y} \right) + \left(\frac{\partial f}{\partial z}F_3 + f\frac{\partial F_3}{\partial z} \right) \\ &= \left(f\frac{\partial F_1}{\partial x} + f\frac{\partial F_2}{\partial y} + f\frac{\partial F_3}{\partial z} \right) + \left(\frac{\partial f}{\partial x}F_1 + \frac{\partial f}{\partial y}F_2 + \frac{\partial f}{\partial z}F_3 \right) \\ &= f \operatorname{Div}(\mathbf{F}) + \mathbf{F} \cdot \operatorname{grad}(f) \end{aligned}$$

2. One parameterization, which is just a variant of the way we parameterize the circle, is to define

$$\begin{aligned} x(t) &= \cos^3(t) \\ y(t) &= \sin^3(t). \end{aligned}$$



Then it's certainly true that $(x(t))^{2/3} + (y(t))^{2/3} = \cos^2(t) + \sin^2(t) = 1$, so this is a parameterization of the curve. Both $\cos^3(t)$ and $\sin^3(t)$ have derivatives of all orders, so this is a C^∞ parameterization. It is therefore also continuous and differentiable. (The curve might not look like it should have such a nice parameterization, but it does.)

3. We want to compute the integral of the function $f(x, y) = xy - x - y + 1 = (x-1)(y-1)$ along different curves connecting $(1, 0)$ and $(0, 1)$.

(a) \mathbf{c}_1 , the circular arc $\mathbf{c}_1(t) = (\cos(t), \sin(t))$, $0 \leq t \leq \pi/2$.

The speed is $\|\mathbf{c}'_1(t)\| = \sqrt{(-\sin(t))^2 + (\cos(t))^2} = 1$, and $f(\mathbf{c}_1(t)) = \cos(t)\sin(t) - \cos(t) - \sin(t) + 1$, so the integral becomes

$$\begin{aligned} \int_{\mathbf{c}_1} f \, ds &= \int_0^{\pi/2} \sin(t)\cos(t) - \cos(t) - \sin(t) + 1 \, dt \\ &= \frac{\sin^2(t)}{2} - \sin(t) + \cos(t) + t \Big|_0^{\pi/2} \\ &= \left(\frac{1}{2} - 1 + 0 + \frac{\pi}{2} \right) - \left(\frac{0}{2} - 0 + 1 + 0 \right) = \frac{\pi-3}{2}. \end{aligned}$$

(b) \mathbf{c}_2 , the straight line segment $\mathbf{c}_2(t) = (1 - t, t)$, $0 \leq t \leq 1$.

The speed is $\|\mathbf{c}'_2(t)\| = \sqrt{(-1)^2 + (1)^2} = \sqrt{2}$, and $f(\mathbf{c}_2(t)) = (1-t)t - (1-t) - t + 1 = t - t^2$, so the integral becomes

$$\begin{aligned} \int_{\mathbf{c}_2} f ds &= \int_0^1 \sqrt{2}(t - t^2) dt \\ &= \sqrt{2} \left(\frac{t^2}{2} - \frac{t^3}{3} \right)_{t=0}^{t=1} = \frac{\sqrt{2}}{6}. \end{aligned}$$

(c) \mathbf{c}_3 , from $(1, 0)$ horizontally to the origin, then vertically to $(0, 1)$.

This curves is parameterized in two pieces. The first piece is $\mathbf{c}_{3,1}(t) = (1 - t, 0)$ for $0 \leq t \leq 1$.

The speed is $\|\mathbf{c}'_{3,1}(t)\| = \sqrt{(-1)^2 + (0)^2} = 1$, and $f(\mathbf{c}_{3,1}(t)) = t$, so the first piece of the integral becomes

$$\int_{\mathbf{c}_{3,1}} f ds = \int_0^1 t dt = \frac{1}{2}.$$

The second piece is $\mathbf{c}_{3,2}(t) = (0, t)$ for $0 \leq t \leq 1$.

The speed is $\|\mathbf{c}'_{3,2}(t)\| = \sqrt{(0)^2 + (1)^2} = 1$, and $f(\mathbf{c}_{3,2}(t)) = 1 - t$, so the second piece of the integral becomes

$$\int_{\mathbf{c}_{3,2}} f ds = \int_0^1 (1 - t) dt = \frac{1}{2}.$$

The total integral along \mathbf{c}_3 is just the sum of these two terms:

$$\int_{\mathbf{c}_3} f ds = \frac{1}{2} + \frac{1}{2} = 1.$$

(d) \mathbf{c}_4 , from $(1, 0)$ vertically to $(1, 1)$, then horizontally to $(0, 1)$.

This is again something we compute in two pieces. The first piece is $\mathbf{c}_{4,1}(t) = (1, t)$ for $0 \leq t \leq 1$.

The speed is $\|\mathbf{c}'_{4,1}(t)\| = \sqrt{(0)^2 + (1)^2} = 1$, and $f(\mathbf{c}_{4,1}(t)) = 0$, so the first piece of the integral is just

$$\int_{\mathbf{c}_{4,1}} f ds = \int_0^1 0 dt = 0.$$

The second piece is $\mathbf{c}_{4,2}(t) = (1 - t, 1)$ for $0 \leq t \leq 1$.

The speed is $\|\mathbf{c}'_{4,2}(t)\| = \sqrt{(-1)^2 + (0)^2} = 1$, and $f(\mathbf{c}_{4,2}(t)) = 0$ again, so again we have

$$\int_{\mathbf{c}_{4,2}} f \, ds = \int_0^1 0 \, dt = 0.$$

The total integral along \mathbf{c}_3 is the sum

$$\int_{\mathbf{c}_4} f \, ds = 0 + 0 = 0.$$

(e) \mathbf{c}_5 , the circular arc $\mathbf{c}_5(t) = (\cos(t), -\sin(t))$, $0 \leq t \leq 3\pi/2$.

The speed is $\|\mathbf{c}'_5(t)\| = \sqrt{(-\sin(t))^2 + (-\cos(t))^2} = 1$, and $f(\mathbf{c}_5(t)) = -\sin(t)\cos(t) - \cos(t) + \sin(t) + 1$, so the integral becomes

$$\begin{aligned} \int_{\mathbf{c}} f \, ds &= \int_0^{3\pi/2} -\sin(t)\cos(t) - \cos(t) + \sin(t) + 1 \, dt \\ &= \left. \frac{\cos^2(t)}{2} - \sin(t) - \cos(t) + t \right|_{t=0}^{t=3\pi/2} \\ &= \left(\frac{0}{2} - (-1) - 0 + \frac{3\pi}{2} \right) - \left(\frac{1}{2} - 0 - 1 + 0 \right) = \frac{3\pi+3}{2}. \end{aligned}$$

4. If \mathbf{c} is the “twisted cubic:” $\mathbf{c}(t) = (t, t^2, t^3)$ for $t \in [0, 4]$, then $\mathbf{c}'(t) = (1, 2t, 3t^2)$, so $\|\mathbf{c}'(t)\| = \sqrt{1 + 4t^2 + 9t^4}$. If the function f is $f(x, y, z) = \sqrt{9xz + 4y + 1}$ then

$$f(\mathbf{c}(t)) = f(t, t^2, t^3) = \sqrt{9 \cdot t \cdot t^3 + 4 \cdot t^2 + 1} = \sqrt{1 + 4t^2 + 9t^4}.$$

Therefore,

$$\begin{aligned} \int_{\mathbf{c}} f \, ds &= \int_0^4 \left(\sqrt{1 + 4t^2 + 9t^4} \right)^2 \, dt \\ &= \int_0^4 1 + 4t^2 + 9t^4 \, dt \\ &= \left. t + \frac{4}{3}t^3 + \frac{9}{5}t^5 \right|_{t=0}^{t=4} = \frac{28988}{15} = 1932 + \frac{8}{15}. \end{aligned}$$

5. To find the average value of a function along a curve, we find its integral along the curve, and divide by the arclength. For the function $f(x, y, z) = xyz$, along the helix

$$\mathbf{c}(t) = (\sin(t), 8t, \cos(t)), \quad t \in [0, 6\pi],$$

the speed is $\|\mathbf{c}(t)\| = \sqrt{(\cos(t))^2 + (8)^2 + (-\sin(t))^2} = \sqrt{65}$, and $f(\mathbf{c}(t)) = 8t \cos(t) \sin(t)$.

Therefore the integral of f along \mathbf{c} is

$$\begin{aligned}\int_{\mathbf{c}} f \, ds &= \int_0^{6\pi} 8\sqrt{65} t \cos(t) \sin(t) \, dt \\ &= \sqrt{65} \left(4t \sin^2(t) + \sin(2t) - 2t \right) \Big|_{t=0}^{t=6\pi} \\ &= \sqrt{65} (0 + 0 - 12\pi) - \sqrt{65} (0 + 0 - 0) = -12\sqrt{65}\pi.\end{aligned}$$

The arclength is easier to find, since the speed is a constant $\sqrt{65}$, the arclength is

$$\int_0^{6\pi} \sqrt{65} \, dt = 6\sqrt{65} \pi.$$

The average value of f along \mathbf{c} is therefore

$$\frac{-12\sqrt{65}\pi}{6\sqrt{65}\pi} = -2.$$