

INVARIANTS OF THE DIAGONAL C_p -ACTION ON V_3 — ADDITIONAL DETAILS

H.E.A. CAMPBELL, B. FODDEN, AND DAVID L. WEHLAU

ABSTRACT. In this document we provide extra details of the proofs and computations given in [0] where a finite SAGBI basis for $\mathbf{F}[V_3 \oplus V_3]^{C_p}$ is given.

The purpose of this document is to provide additional details to the interested reader of [0]. These details are omitted from [0] for reasons of clarity and brevity. This document is not intended to stand on its own and we assume the reader has already read [0]. All the notation and definitions used here are given in [0].

1. LEAD MONOMIALS OF TRANSFERS

Here we prove Lemma 5.1 which is needed to identify the lead monomials of some of the transfers lying in the set B . This lemma is a simple extension of the results found in [14, § 3].

It is easily seen that for $j = 1, 2$ and $0 \leq c \leq p - 1$

$$\begin{aligned}\sigma^c(z_j) &= z_j + cy_j + \binom{c}{2}x_j \\ \sigma^c(y_j) &= y_j + cx_j.\end{aligned}$$

Furthermore it is well known (for a proof see for example [5, 9.4]) that

$$\sum_{c=0}^{p-1} c^i = \begin{cases} 0 \text{ in } \mathbf{F} & \text{if } i < p - 1 \\ -1 \text{ in } \mathbf{F} & \text{if } i = p - 1 \end{cases}$$

In particular, if $f(c)$ is any polynomial of degree less than $p - 1$ then $\sum_{c=0}^{p-1} f(c) = 0$ in \mathbf{F} .

Lemma 5.1. *Suppose $0 \leq t, s \leq p - 1$ and $2s + t \geq p - 1$. Then*

$$\text{LM}(\text{Tr}(z_1^t z_2^s)) = \begin{cases} z_1^t x_2^{p-1-s} y_2^{2s-(p-1)} & \text{if } s \geq (p-1)/2 \\ y_1^{p-1-2s} z_1^{t+2s-(p-1)} x_2^s & \text{if } s \leq (p-1)/2 \end{cases}$$

Proof. If $s \geq (p-1)/2$ then this result is a minor modification of [14, Theorem 3.2]. Therefore we suppose that $s \leq (p-1)/2$. We have $\sigma^c(z_1^t z_2^s) = (z_1 + cy_1 + \binom{c}{2}x_1)^t (z_2 + cy_2 + \binom{c}{2}x_2)^s$. Expanding this we

Date: January 20, 2005.

1991 Mathematics Subject Classification. 13A50.

Research partially supported by grants from ARP and NSERC.

obtain an expression $\text{Tr}(z_1^t z_2^s) = \sum_{c=0}^{p-1} \sum_w f_w(c) \mathbf{x}^w$ where $f_w \in \mathbf{F}[c]$ for each w . The largest monomial w here (with respect to the monomial ordering) for which the polynomial f_w has degree at least $p-1$ is the monomial $\mathbf{x}^w = y_1^{p-1-2s} z_1^{t+2s-(p-1)} x_2^s$. By the above lemma, all larger monomials have coefficient 0 in $\text{Tr}(z_1^t z_2^s)$. For $w = y_1^{p-1-2s} z_1^{t+2s-(p-1)} x_2^s$ we get $f_w(c) = \binom{t}{p-1-2s} c^{p-1-2s} \binom{c}{2}^{2s} = 2^{-2s} c^{p-1} \left(\binom{t}{p-1-2s} - \binom{t}{p-1-2s} c^{-s} \right)$ and thus $\sum_{c=0}^{p-1} f_w(c) = -\binom{t}{p-1-2s} 2^{-2s} \neq 0$ since $0 \leq p-1-2s \leq t \leq p-1$. Therefore $\text{LT}(\text{Tr}(z_1^t z_2^s)) = -2^{-2s} \binom{t}{p-1-2s} y_1^{p-1-2s} z_1^{t+2s-(p-1)} x_2^s$. \square

Applying Lemma 5.1 together with simple generalizations of [14, Theorem 3.3] and [14, Theorem 3.6] we find that M is generated over A by the following monomials.

- (0) 1
- (1) $z_1^s x_2^s$ for $1 \leq s \leq (p-3)/2$
- (2) $y_1 z_1^s x_2^{s+1}$ for $0 \leq s \leq (p-1)/2$
- (3a) $y_1^{p-2s} z_1^{2s} x_2^s$ for $0 \leq s \leq (p-1)/2$
- (3b) $y_1 z_1^{p-1} x_2^{p-1-s} y_2^{2s-(p-1)}$ for $(p+1)/2 \leq s \leq p-1$
- (4) $z_1^s y_2^p$ for $0 \leq s \leq p-1$
- (5a) $y_1^{p-1-2s} z_1^{t+2s-(p-1)} x_2^s$ for $1 \leq s \leq (p-1)/2, (p+1)/2 \leq t \leq p-1$
and $p \leq t+s$
- (5b) $z_1^t x_2^{p-1-s} y_2^{2s-(p-1)}$ for $(p+1)/2 \leq s \leq p-1, 1 \leq t \leq p-1$ and
 $p \leq t+s$
- (6a) $y_1^{p-2s} z_1^{t+2s-(p-1)} x_2^s$ for $1 \leq s \leq (p-1)/2, (p-1)/2 \leq t \leq p-2$
and $p-1 \leq t+s$
- (6b) $y_1 z_1^t x_2^{p-1-s} y_2^{2s-(p-1)}$ for $(p+1)/2 \leq s \leq p-1, 0 \leq t \leq p-2$ and
 $p-1 \leq t+s$
- (7) $y_1 z_1^s y_2^p$ for $0 \leq s \leq p-1$

The numbering of these families of lead monomials corresponds to the numbering of the families of invariants given in Theorem 4.1.

2. COMPUTATION OF $\mathcal{H}(M, \lambda)$

In this section we give addition details of the computation of the Hilbert series of M .

Decompose M by multi-degree (with respect to the H -grading) as follows:

$$M = \bigoplus_{\omega \in C_2 \times C_2 \times C_p} M_\omega = \bigoplus_{i=0}^1 \bigoplus_{j=0}^1 \bigoplus_{k=0}^{p-1} M_{(i,j,k)}.$$

By this direct sum decomposition,

$$\mathcal{H}(M, \lambda) = \sum_{i=0}^1 \sum_{j=0}^1 \sum_{k=0}^{p-1} \mathcal{H}(M_{(i,j,k)}, \lambda)$$

and we compute $\mathcal{H}(M, \lambda)$ by computing each of the individual $\mathcal{H}(M_{(i,j,k)}, \lambda)$.

To do this we begin by sorting the monomials in C according to their H -degree.

For $i = 0$ and all $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$, if $k + j \leq (p - 1)/2$ we have the following generators of $M_{(0,j,k)}$.

- (1) $y_1^{p-1+j-2t} z_1^k x_2^t$ for $\lceil k/2 \rceil \leq t \leq k + j - 1$.
- (2) $y_1^j z_1^k x_2^{k+j}$
- (3) $y_1^j z_1^k x_2^{p-1-t} y_2^{2t-(p-1)}$ for $p - j - k \leq t \leq p - 1$

For $i = 0$ and all $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$ if $k + j > (p - 1)/2$ we have the following generators of $M_{(0,j,k)}$.

- (1) $y_1^{p-1+j-2t} z_1^k x_2^t$ for $\lceil k/2 \rceil \leq t \leq (p - 1)/2$.
- (2) $y_1^j z_1^k x_2^{p-1-t} y_2^{2t-(p-1)}$ for $(p + 1)/2 \leq t \leq p - 1$

For $i = 1$, and all $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$ we have the following generator of $M_{(1,j,k)}$.

- (1) $y_1^j z_1^k y_2^p$

Since each $M_{(1,j,k)}$ is generated over A by a single element, it is a free rank one A -module. Therefore

$$(2.0.1) \quad \mathcal{H}(M_{(1,j,k)}, \lambda) = \mathcal{H}(A, \lambda) \cdot \lambda^{p+k+j}$$

for all $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$.

In Equations (5.1.3) and (5.1.4) we give expressions for $\mathcal{H}(M_{(0,j,k)}, \lambda)$. Here is an expanded description of the derivation of these expressions.

Examining the points in $V_{(0,j,k)}$ we find the following values for the $\text{LCM}(\Delta_1)$ and the $\text{LCM}(\Delta_2)$.

For $i = 0$ and $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$, if $j + k \leq (p - 1)/2$ then we have the following points in $\{\text{LCM}(\Delta_1) \mid \Delta_1 \in E_{(0,j,k)}\}$.

- (1) $(0, j, k, p - t, 2t - (p - 1), 0)$ for $p - j - k \leq t \leq p - 1$
- (2) $(0, p + j - 1 - 2t, k, t + 1, 0, 0)$ for $\lceil k/2 \rceil \leq t \leq k + j - 1$
- (3) $(0, p + j - 1 - 2t, k, t, p - 1 - 2t, 0)$ for $\lceil k/2 \rceil \leq t \leq k + j - 1$

For $i = 0$ and $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$, if $j + k \geq (p + 1)/2$ then we have the following points in $\{\text{LCM}(\Delta_1) \mid \Delta_1 \in E_{(0,j,k)}\}$.

- (1) $(0, j, k, p - t, 2t - (p - 1), 0)$ for $(p + 1)/2 \leq t \leq p - 1$
- (2) $(0, p + j - 1 - 2t, k, t + 1, 0, 0)$ for $\lceil k/2 \rceil \leq t \leq (p - 3)/2$
- (3) $(0, p + j - 1 - 2t, k, t, p - 1 - 2t, 0)$ for $\lceil k/2 \rceil \leq t \leq (p - 3)/2$

For $i = 0$ and $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$, if $j + k \leq (p - 1)/2$ then we have the following points in $\{\text{LCM}(\Delta_2) \mid \Delta_2 \in F_{(0,j,k)}\}$.

- (1) $(0, p + j - 1 - 2t, k, t + 1, p - 1 - 2t, 0)$ for $\lceil k/2 \rceil \leq t \leq k + j - 1$

For $i = 0$ and $0 \leq j \leq 1$ and $0 \leq k \leq p - 1$, if $j + k \geq (p + 1)/2$ then we have the following points in $\{\text{LCM}(\Delta_2) \mid \Delta_2 \in F_{(0,j,k)}\}$.

- (1) $(0, p + j - 1 - 2t, k, t + 1, p - 1 - 2t, 0)$ for $\lceil k/2 \rceil \leq t \leq (p - 3)/2$

Therefore if $j + k \leq (p - 1)/2$ then

$$\begin{aligned}\mathcal{H}(K_0, \lambda) &= \left(\lambda^{2k+2j} + \sum_{t=\lceil k/2 \rceil}^{k+j-1} \lambda^{p-1+j+k-t} + \sum_{t=p-j-k}^{p-1} \lambda^{j+k+t} \right) \mathcal{H}(A, \lambda) \\ \mathcal{H}(K_1, \lambda) &= \left(\sum_{t=p-j-k}^{p-1} \lambda^{j+k+t+1} + \sum_{t=\lceil k/2 \rceil}^{k+j-1} (\lambda^{p+j+k-t} + \lambda^{2p+j+k-3t-2}) \right) \mathcal{H}(A, \lambda) \\ \mathcal{H}(K_2, \lambda) &= \left(\sum_{t=\lceil k/2 \rceil}^{k+j-1} \lambda^{2p+j+k-3t-1} \right) \mathcal{H}(A, \lambda)\end{aligned}$$

Conversely if $k + j \geq (p + 1)/2$ then

$$\begin{aligned}\mathcal{H}(K_0, \lambda) &= \left(\sum_{t=\lceil k/2 \rceil}^{(p-1)/2} \lambda^{p-1+j+k-t} + \sum_{t=(p+1)/2}^{p-1} \lambda^{j+k+t} \right) \mathcal{H}(A, \lambda) \\ \mathcal{H}(K_1, \lambda) &= \left(\sum_{t=(p+1)/2}^{p-1} \lambda^{j+k+t+1} + \sum_{t=\lceil k/2 \rceil}^{(p-3)/2} (\lambda^{p+j+k-t} + \lambda^{2p+j+k-3t-2}) \right) \mathcal{H}(A, \lambda) \\ \mathcal{H}(K_2, \lambda) &= \left(\sum_{t=\lceil k/2 \rceil}^{(p-3)/2} \lambda^{2p+j+k-3t-1} \right) \mathcal{H}(A, \lambda)\end{aligned}$$

From the above, using (5.1.2) and observing that the resulting sums telescope, we obtain the expressions given in (5.1.3) and (5.1.4).

REFERENCES

- [0] H E A Campbell, B Fodden, and David L Wehlau, *Invariants of the Diagonal C_p -action on V_3* , J. Algebra, ? No ? (2005?) ?-?.
- [5] H E A Campbell, I P Hughes, R J Shank, and D L Wehlau, *Bases for rings of covariants*, Transformation Groups, **1** No 4 (1996) 307–336.
- [14] R. James Shank, *S.A.G.B.I. bases for rings of formal modular seminvariants*, Comment. Math. Helv. **73** (1998) 548–565.

DEPARTMENT OF MATHEMATICS & STATISTICS, MEMORIAL UNIVERSITY OF
NEWFOUNDLAND, ST JOHN'S NL, CANADA A1C 5S7
E-mail address: vpacad@mun.ca

DEPARTMENT OF MATHEMATICS & STATISTICS, QUEEN'S UNIVERSITY, KINGSTON,
ONTARIO, CANADA K7L 3N6
E-mail address: fodden@mast.queensu.ca

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE,
ROYAL MILITARY COLLEGE, KINGSTON, ONTARIO, CANADA K7K 7B4
E-mail address: wehlau@rmc.ca