

# Hurwitz Spaces of Genus 2 Covers of an Elliptic Curve

## 1. Introduction

**Hurwitz spaces:** – introduced by Hurwitz in 1891.

– classify covers  $f : Y \rightarrow X$  of degree  $N$  over a fixed curve  $X/K$  with  $w$  ramification points  $R_f \subset X$ .

**Problem (Hurwitz)** Fix  $X/K$ ,  $N$ ,  $w$  and  $R \subset X$ .

1) Investigate the totality  $H(X/K, N, w)$  of all covers  $f : Y \rightarrow X$  with  $\deg(f) = N$  and  $\#R_f = w$ .

2) Calculate the number  $\#H(X/K, N, R)$  of such covers with  $R_f = R$ .

**Remarks:** 1) Throughout, we always consider equiv-

alence classes of covers:  $(Y_1 \xrightarrow{f_1} X) \sim (Y_2 \xrightarrow{f_2} X) \Leftrightarrow \exists \phi : Y_1 \xrightarrow{\sim} Y_2$  with  $f_2 \circ \phi = f_1$ .

2) A cover  $f : Y \rightarrow X$  is called simple if

$$f^{-1}(x) \geq \deg(f) - 1, \quad \text{for all } x \in X.$$

Such covers constitute an important subclass of all covers.

**Theorem A (Hurwitz, 1891):** If  $K = \mathbb{C}$ , then

- (a)  $H(X/\mathbb{C}, N, w)$  is a “Riemannian space”.
- (b)  $H^{simple}(\mathbb{P}^1/\mathbb{C}, N, w)$  is a **connected** manifold of dimension  $w$  (provided that  $w \geq 2N - 2$  and  $2|w$ ).
- (c) The **discriminant map**

$$\delta : H^{simple}(\mathbb{P}^1/\mathbb{C}, N, w) \rightarrow (\mathbb{P}^1)^{(w)} \setminus \Delta_w$$

is **finite** and **etale**. Thus,  $\#H^{simple}(\mathbb{P}^1/\mathbb{C}, N, R)$  depends only on  $w = \#R$ .

**Observation (Hurwitz):** RET  $\Rightarrow$  the calculation of  $\#H(X/\mathbb{C}, N, R)$  is a **purely group-theoretic problem**, albeit one that is “highly complicated” (**Hurwitz**).

**Hurwitz (1891/1901)** found a “satisfactory solution” for calculating  $n_{N,w} := \#H^{simple}(\mathbb{P}^1/\mathbb{C}, N, R)$ :

$$\begin{aligned} n_{3,w} &= \frac{1}{3!}(3^{w-1} - 3), & (n_{2,w} = 1) \\ n_{4,w} &= \frac{1}{4!}(2^{w-2} - 4)(3^{w-1} - 3), \text{ etc.} \end{aligned}$$

**Question:** What about **other** ground fields?

**Hurwitz (1891):** partial results for  $K = \mathbb{R}$ .

**Fulton (1969):** consider covers of **curve families**/ $S$ :

$$f : \mathcal{Y} \rightarrow \mathbb{P}_S^1 = \mathbb{P}^1 \times S, \quad S \text{ any scheme.}$$

**Thus:** 1) The fibres  $f_s : \mathcal{Y}_s \rightarrow (\mathbb{P}_S^1)_s = \mathbb{P}_{\kappa(s)}^1$  of  $f$  are covers of curves in the previous sense.

2) The assignment  $S \mapsto H^{simple}(\mathbb{P}_S^1/S, N, w)$  defines a **functor**  $\mathcal{H}_{N,w} : \underline{Sch} \rightarrow \underline{Sets}$ .

**Theorem C (Fulton, 1969):** If  $N \geq 3$ , then the functor  $\mathcal{H}_{N,w}$  is **(finely) representable** by a scheme  $H_{N,w}/\mathbb{Z}$  of finite type. In particular, for any field  $K$  we have  $H_{N,w}(K) = H^{simple}(\mathbb{P}^1/K, N, w)$ .

In addition, the restriction of the **discriminant map** to  $H_{N,w} \otimes \mathbb{Z}[1/N!] \subset H_{N,w}$ ,

$$\delta : H_{N,w} \otimes \mathbb{Z}[1/N!] \rightarrow (\mathbb{P}_{\mathbb{Z}[1/N!]}^1)^{(w)} \setminus \Delta_w,$$

is **finite** and **etale**.

**Remark:** Little seems to be known about the **geometric structure** of  $H_{N,w}$ .

**Aim:** Study **analogues** of these results in the case that  $X = E$  is an **elliptic curve** (and  $w = 2$ ).

## 2. The Case $X = E$ and $w = 2$ ( $\Rightarrow g_Y = 2$ ).

**Reference:** IEM Preprint No. 9 (2001), IEM Essen.

(See also [www.mast.queensu.ca/~kani](http://www.mast.queensu.ca/~kani).)

**Let**  $E/K$  be an elliptic curve over a field  $K$  ( $\text{char} \neq 2$ ).

1)  $E(K)$  acts on  $E$  and hence on  $H(E/K, N, 2)$  via translation:  $f \mapsto T_x \circ f$ . Thus  $H \sim H' \times E$ .

2) Each  $Y \xrightarrow{f} E$  factors as  $Y \xrightarrow{f'} E' \xrightarrow{u_f} E$ , where  $u_f : E' \rightarrow E$  is the max. unramified subcover of  $f$ .

Thus:  $H(\dots)$  is a union of components which are indexed by subgroups  $G \leq E$  with  $\#G | N$  (and  $\#G \neq N$ ); explicitly,  $G = \text{Ker}(\hat{u}_f)$ .

**Definition:** a)  $f$  is called minimal if  $\deg(u_f) = 1$ .

b) A cover  $f : Y \rightarrow E$  with  $g_Y = 2$  is called normalized if it is minimal and if

$$f(W) \subset E[2] \text{ and } \#(f^{-1}(0_E) \cap W) = \begin{cases} 3 & N \text{ odd} \\ 0 & \text{else} \end{cases}$$

where  $W = \text{Fix}(\sigma_Y)$  denotes the set of 6 Weierstrass points of  $Y$ . (Here:  $\sigma_Y$  is the hyperelliptic involution of  $Y$ .)

**Notes:** 1) If  $f : Y \rightarrow E$  is **minimal**, then  $\exists! x \in E(K)$  such that  $T_x \circ f : Y \rightarrow E$  is **normalized**.

2) If  $f$  is **normalized**, then  $f \circ \sigma_Y = [-1]_E \circ f$ . Thus  $\text{Disc}(f)$  is **symmetric** with respect to  $[-1]_E$ , i.e.  $[-1]_E^* \text{Disc}(f) = \text{Disc}(f)$ .

**Example:** Let

$$E : y^2 = (x - a)(x - b)(x - c), \quad abc \neq 0$$

$$Y : s^2 = (t^2 - a)(t^2 - b)(t^2 - c).$$

Then the cover  $f : Y \rightarrow E$ , given by  $f^*x = t^2$ ,  $f^*y = s$ , is **normalized** and of degree 2.

**Remark:** The notion of **normalized curve covers** can be extended naturally to **families of curve covers**  $f : \mathcal{Y} \rightarrow E_S = E \times S$  so as to obtain a functor

$$\mathcal{H} = \mathcal{H}_{E/K, N} : \underline{Sch}/K \rightarrow \underline{Sets}$$

given by  $S \mapsto \mathcal{H}(S) = \{\mathcal{Y} \xrightarrow{f} E_S : f \text{ is normalized, } \deg(f) = N\}$ .

More generally, if  $E/S$  is a **family** of elliptic curves over a **base**  $S$ , then one obtains in a similar way a functor

$$\mathcal{H}_{E/S, N} : \underline{Sch}/S \rightarrow \underline{Sets}.$$

**Theorem 1:** If  $N \geq 3$  and  $\text{char}(K) \nmid N$ , then the functor  $\mathcal{H}_{E/K,N}$  is finely represented by a smooth, affine and geometrically connected curve  $H_{E/K,N}/K$ . More precisely,  $H_{E/K,N}$  is an open subscheme of a certain twist  $X_{E/K,N,-1}$  of the modular curve  $X(N)$  of level  $N$ . In particular,

$$H_{E/K,N} \otimes \overline{K} \stackrel{\text{open}}{\subset} X'(N)_{/\overline{K}} \subset X(N)_{/\overline{K}}.$$

**Remarks:** 1) If  $K = \mathbb{C}$ , then  $X'(N) = \Gamma(N) \backslash \mathfrak{H}$  and  $X(N) = \Gamma(N) \backslash \mathfrak{H}^*$ , which is a Galois cover of  $X(1) \simeq \mathbb{P}^1$  of degree

$$\overline{sl}(N) := |\text{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm 1\}|.$$

2) Theorem 1 extends to families of elliptic curves  $E/S$ , where  $S$  any scheme (with  $\frac{1}{2N} \in S$ ).

**Example:** Let  $S = X'(N)_{/\mathbb{Z}[\zeta_N, 1/N]}$  and  $E = E(N)$ , the universal elliptic curve over  $X'(N)$  (with level  $N$  structure). Then  $\mathcal{H}_{E(N)/X'(N),N}$  is represented by an open affine subscheme

$$H_{E(N)/X'(N),N} \subset X'(N) \times X'(N).$$

**Theorem 2:** Let

$$D_{E/K,N} = X(N)_{/\overline{K}} \setminus (H_{E/K,N} \otimes \overline{K})$$

denote the **degeneracy locus**. Then

$$\#D_{E/K,N} \leq \frac{1}{12N}(5N + 6)\overline{sl}(N),$$

and equality holds if and only if  $\text{char}(K) \nmid N!$ .

– **reinterpretation** of results of **Crelle J. 485 (1997)**  
+ **J. No. Th. 64 (1997)**.

**Theorem 3:** The assignment  $(Y \xrightarrow{f} E) \mapsto \text{Disc}(f)$  is represented by a **quasi-finite** morphism

$$\delta = \delta_{E/K,N} : H_{E/K,N} \rightarrow \mathbb{P}_K^1 \simeq (E^{(2)})^{sym}.$$

Furthermore, if  $\text{char}(K) \nmid N!$ , then  $\delta$  is **finite** and **etale** outside of  $\pi_E(E[2]) \subset \mathbb{P}^1$ .

**Theorem 4:** If  $\text{char}(K) \nmid N!$ , then

$$\deg(\delta_{E/K,N}) = \frac{1}{6}(N - 1)\overline{sl}(N).$$

**Remarks:** 1) This degree can be viewed as a **measure of non-rigidity** of coverings ( $\rightarrow$  **Völklein**).

2) **H. Völklein** proved **Theorem 4** for  $N = 3, 5, 7$  by using **group theory** (and a computer).

### 3. Some applications

#### (a) Rationality Questions ( $K$ a number field)

Since  $g_{X(N)} \geq 2$  for  $N \geq 7$ , we have by **Faltings' theorem** (= **Mordell's Conjecture**):

**Corollary 1:**  $\#\mathcal{H}_{E/K,N}(K) < \infty$ , if  $N \geq 7$ .

**Question:** Is  $\mathcal{H}_{E/K,N}(K) = \emptyset$ , for  $N \gg 0$ ?

This is **false** (even for  $N$  **prime**), for there exist curves  $Y/K$  with  $\infty$ 'ly many  $f_N : Y \rightarrow E$  for which  $N = \deg(f_N)$  is **prime**.

**Conjecture (\*)** For each  $E/K$  there exist only **finitely many** genus 2 curves  $Y/K$  which have a (minimal) morphism  $f : Y \rightarrow E$  of degree  $N \geq 7$ .

**Remark:** **ABC conj.**  $\Rightarrow$  **Asym. Fermat**  $\Rightarrow$  **Conj. (\*)**.

Moreover, the converse: **Conj. (\*)**  $\Rightarrow$  **Asym. Fermat** is “almost true”: it implies a slightly weaker version of **Frey's Conjecture 5** (which by **Frey** and **Wiles** is equivalent to the **Asymptotic Fermat Conjecture** (for  $K = \mathbb{Q}$ ).)



## (b) Moduli

**Question:** For which curves  $Y/K$  does there exist a (minimal) morphism  $f : Y \rightarrow E$  of degree  $N$ ?

**Corollary 2:** For every  $N$  there exists a morphism

$$\mu_N : H_{E(N)/X'(N),N} \rightarrow M_2$$

to the **moduli space** of curves of genus 2. Moreover:

- a)  $\text{Im}(\mu_N) = \text{Humbert surface}$  with  $\text{Inv. } \Delta = N^2$ ;
- b)  $\text{deg}(\mu_N) = 2\overline{sl}(N)$ ; more precisely,

$$\text{Im}(\mu_N) \sim Z_{N,-1}^{sym} := (X(N) \times X(N)) / \langle \Delta_{N,-1}, \tau \rangle,$$

where  $\tau(x, y) = (y, x)$  and

$$\Delta_{N,-1} = \{(g, \alpha_{-1}(g)) : g \in \text{SL}_2(\mathbb{Z}/N\mathbb{Z}) / \{\pm 1\}\},$$

where  $\alpha_{-1}(g) = Q_{-1}gQ_{-1}^{-1}$  with  $Q_{-1} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ .

**In particular,** the **normalization** (and compactification) of the **Humbert surface**  $\text{Im}(\mu_N)$  is the **symmetric Diagonal Quotient Surface**  $Z_{N,-1}^{sym}$ .

### (c) Counting Covers: ( $K = \overline{K}$ )

**Corollary 3:** If  $N \geq 2$  and  $\text{char}(K) \nmid N$ , then for every  $R \subset E$  with  $\#R = 2$  we have

$$c_N := \sum_{f \in H^s(E/K, N, R)} \frac{1}{|\text{Aut}(f)|} = \frac{1}{3}(N\sigma_3(N) - N^2\sigma_1(N)),$$

where  $\sigma_k(N) = \sum_{d|N} d^k$ . Thus, if  $\text{char}(K) = 0$ , then  $F_2(q) := \sum c_N q^N$  is a **quasi-modular form** of weight 6; explicitly we have

$$(1) \quad F_2(q) = \frac{1}{51840}(10E_2^3 - 6E_2E_4 - 4E_6),$$

where  $E_k = 1 + b_k \sum_{n \geq 1} \sigma_{k-1}(n)q^n$  with  $b_2 = -24$ ,  $b_4 = 240$  and  $b_6 = -504$ .

**Remarks:** 1) The identity (1) was first proven by R. Dijkgraaf (1995) by using the methods of **mirror symmetry**.

2) **Theorem 4**  $\Rightarrow$  **Corollary 3** by using the identities

$$\sum_{n|N} \sigma_1(n) \text{sl}(N/n) = \sigma_3(N),$$

$$\sum_{n|N} n \sigma_1(n) \text{sl}(N/n) = N^2 \sigma_1(N).$$

## (d) Curves with minimal degeneration:

**Let**  $H = H_{E/K,N} \subset X = X(N)$  be the moduli space,  
 $f : Y_N \rightarrow E_H = E \times H$  the universal cover,  
 $p : \bar{Y}_N \rightarrow X$  the minimal model of  $Y_N$  over  $X$ ,  
 $h_{\bar{Y}_N/X} = \deg_X(p_*\omega_{Y_N/X}^0)$  its modular height.

**Corollary 4:** The curve  $\bar{Y}_N/X(N)$  is semi-stable and has bad reduction at  $X \setminus H$ . Furthermore, its Jacobian  $J = J_N$  has bad reduction at  $X(N)_\infty := X(N) \setminus X'(N)$ , and its modular height is

$$h_{\bar{Y}_N/X} = h_{J/X} = \frac{1}{2}(2g_{X(N)} - 2 + \#X(N)_\infty).$$

In particular, for  $N = 3$  one thus obtains a semi-stable family  $p : \bar{Y}_3 \rightarrow \mathbb{P}^1$  whose Jacobian has precisely 4 places of bad reduction.

**Remarks:** 1) By a theorem of Faltings it follows (in char = 0) that for any such curve we have the inequality

$$h_{\bar{Y}_N/X} = h_{J/X} \leq \frac{1}{2}(2g_{X(N)} - 2 + \#X(N)_\infty).$$

2) In a recent preprint E. Viehweg and K. Zuo study the structure of families of abelian varieties with such “minimal degeneration”.

## 4. The Basic Construction

**Reference:** Frey/K., Curves of genus 2 covering elliptic curves ... (Texel Conference, 1989)

**Given:**

$$\begin{array}{ccc}
 Y & & Y \\
 f \downarrow \rightsquigarrow & \swarrow & \searrow \\
 E & & E^\perp
 \end{array}
 \rightsquigarrow \psi : E[N] \xrightarrow{\sim} E^\perp[N].$$

(via the duality theory of  $J_Y$ .)

**Conversely:** given anti-isometry  $\psi : E[N] \rightarrow E'[N]$ , one can recover a (normalized) genus 2 cover

$$f_\psi : Y_\psi \rightarrow E.$$

**However:** the curve  $Y_\psi$  may be reducible!

$$\Rightarrow H_{E/K,N} \subset X_{E/K,N,-1}.$$

**Note:** 1) The moduli space  $X_{E/K,N,-1}$  classifies pairs  $(E', \psi)$ , where  $\psi : E[N] \rightarrow E'[N]$  is an anti-isometry.  
 2) This construction also works for families! (Cf. IEM Preprint, op. cit.):  $\Rightarrow$  Theorem 1.

## 5. Study of Degenerations

**Let**  $H = H_{E/\overline{K}, N}$  denote the moduli space,  
 $f_H : Y_H \rightarrow E_H = E \times_{\overline{K}} H$  the universal cover,  
 $X = X(N) \supset H$  the natural compactification,  
 $\overline{Y}/X$  the minimal model of the generic fibre of  $Y_H$ .

**Facts.** 1) The fibres of  $\overline{Y}/X$  are semi-stable.

2)  $f_H$  extends to a morphism  $f = f_X : \overline{Y} \rightarrow E_X$  which is finite if and only if  $\text{char}(K) \nmid N!$ .

**Theorem 5:** Suppose  $\text{char}(K) \nmid N!$ . Then:

(a) The fibres  $\overline{Y}_x$  of  $\overline{Y}/X$  are stable curves with at most one singularity.

(b)  $\overline{Y}_x$  is singular if and only if  $x \in D_{E/\overline{K}, N} = X_\infty \dot{\cup} X_1$ , where  $X_\infty$  is the set of cusps of  $X$ . (Note that  $\#X_\infty = \overline{sl}(N)/N$ .)

(c) If  $x \in X_\infty$ , then  $\overline{Y}_x$  is an irreducible curve whose normalization is a curve of genus 1.

(d) If  $x \in X_1$ , then  $\overline{Y}_x = E_{x,1} \cup E_{x,2}$  is the union of two curves of genus 1 which meet transversely in a unique point  $P_x$ .

## 6. Calculation of Intersection Numbers

**Let**  $F = \kappa(X)$  denote the function field of  $X = X(N)$ ,  
 $f_F : Y_F \rightarrow E_F$  the **generic cover** over  $F$ ,  
 $D_F = \text{Diff}(f_F)$  the **different divisor** of  $f_F$ ,  
 $W_{C_F} \in \text{Div}(Y_F)$  the **hyperelliptic divisor** of  $Y_F$ ,  
 $D$  and  $W$  their respective **closures** in  $\bar{Y}$ ,  
 $\omega_{\bar{Y}/X}^0$  the **relative dualizing sheaf** of  $p_{\bar{Y}} : \bar{Y} \rightarrow X$ .

**Theorem 6:** The **modular height** of  $\bar{Y}/X$  is

$$h_{\bar{Y}/X} := \deg((p_{\bar{Y}})_*(\omega_{\bar{Y}/X}^0)) = \frac{1}{12}\overline{sl}(N),$$

and the **self-intersection number** of  $\omega_{\bar{Y}/X}^0$  is

$$(\omega_{\bar{Y}/X}^0)^2 = \frac{7}{5}\#X_1 + \frac{1}{5}\#X_\infty = \frac{1}{12N}(7N - 6)\overline{sl}(N).$$

**Remark:** The proof uses **Theorem 2**, the **Noether formula** and **Mumford's formula** (which holds if  $g = 2$ ):

$$h = \omega^2 + \delta_0 + \delta_1 \quad \text{and} \quad 5\omega^2 = \delta_0 + 7\delta_1,$$

where  $h = h_{\bar{Y}/X}$ ,  $\omega = \omega_{\bar{Y}/X}^0$ , and  $\delta_0$  (respect.  $\delta_1$ ) is the number of **singular points** of all fibres which **do not** (respect. **do**) **disconnect** the fibre.

**Theorem 7:** (a)  $D$  is an **irreducible** curve on  $\bar{Y}$  which represents the **dualizing sheaf**:  $\omega_{\bar{Y}/X}^0 \sim D$ .

(b) If  $q_1 = pr_1 \circ f|_D : D \rightarrow E$  and  $q_2 = pr_2 \circ f|_D : D \rightarrow X$ , then  $\pi_E \circ q_1 = \bar{\delta}_{E,N} \circ q_2$ , where  $\bar{\delta} : X \rightarrow \mathbb{P}^1$  is the unique extension of  $\delta : H \rightarrow \mathbb{P}^1$ . Thus

$$\deg(\bar{\delta}) = \deg(q_1) = (\omega_{\bar{Y}/X}^0 \cdot f^*(P \times X)).$$

(c) We have  $6D \sim 2W + f^*(E \times A)$ , for some  $A \in \text{Div}(X)$ , and hence

$$\deg(q_1) = \frac{N}{6} \deg(A) = \frac{N}{36} (9(\omega_{\bar{Y}/X}^0)^2 - W^2).$$

(d) The self-intersection number of  $W$  is

$$W^2 = \frac{6}{7} \#X_1 - \frac{9}{7} \omega^2 = -\frac{3}{4N} (N-2) \bar{s}l(N).$$

**Remark:** To compute  $W^2$ , consider the **pullback**  $W^*$  of  $W$  to (the desingularization of)  $\bar{Y} \times_X X(2N)$ , and observe that  $W^* = W_1 + \dots + W_6 + B$ , where the  $W_i$ 's are **6 disjoint sections** and  $B$  is a **fibrar divisor** supported on the fibres over  $X(2N)_\infty$ .